

Interaction manifolds for reaction diffusion equations in 2D

J. Douglas Wright
Department of Mathematics
Drexel University
3141 Chestnut St
Philadelphia, PA 19104
jdoug@math.drexel.edu

September 16, 2009

Abstract

We consider a general planar reaction diffusion equation which we hypothesize has a localized traveling wave solution. Under assumptions which are no stronger than those needed to prove the stability of a single pulse, we prove that the PDE has solutions which are roughly the linear superposition of two pulses, so long as they move along trajectories which are not parallel. In particular we prove that if the initial data for the equation is close to the sum of two separated pulses, then the solution converges exponentially fast to such a superposition so long as the distance between the two pulses remains sufficiently large.

1 Introduction

1.1 The system and hypotheses

Given the existence and stability of localized traveling wave solutions (a.k.a. pulses) to a reaction diffusion equation, one expects that there are solutions (called *multipulse* solutions) which are nearly the linear superposition of two or more such pulses, at least until such time as those pulse come close to one another. There are rigorous justifications for this conjecture in a number of settings, most of which are posed for one spatial variable. [1, 7, 8, 12, 18] treat existence and stability of multipulse standing solutions and use primarily dynamical systems and geometric techniques. [5, 9, 10, 11, 14, 22] handle counter-propagating fronts and pulses and give very complete descriptions of interactions by means of the maximum principle. In [6, 23], the authors handle long distance weak interactions between standing pulses—in particular they compute the induced motion between the pulses. [23] is particularly notable here in that it covers any number of pulses in any spatial dimension. Our work here is most closely related to that of [4, 19, 21], which deal with counter-propagating fronts or pulses in reaction diffusion systems. The chief difficulty in this case is the fact that the problem cannot be made autonomous by transforming into a moving reference frame.

For planar reaction diffusion systems, a host of numerical simulations bear out the expectation (see [15, 17]) that there are solutions which are roughly the sum of pulses moving in different directions and in this paper we prove the corresponding rigorous results. Consider:

$$u_t = \mathcal{D}\Delta u + F(u) \tag{1}$$

where

$$\mathbf{x} = (x_1, x_2) \in \mathbf{R}^2, \quad u(\mathbf{x}, t) \in \mathbf{R}^N$$

and

$$F \in C^2, \quad \mathcal{D} \text{ is a diagonal, positive matrix.}$$

Equations of this type model gas-discharge systems, excitable media in cardiac tissue, numerous chemical reactions, cell growth, blood clotting, *etc.* We make precise our assumptions about existence and stability of pulses.

Hypothesis 1. *There exists a C^3 function $Q(\mathbf{y})$ and $c > 0$ such that*

$$u(\mathbf{x}, t) = Q(\mathbf{x} - ct\mathbf{i})$$

solves (1). Here $\mathbf{i} = (1, 0)$. Furthermore, we assume there exists $\beta_1 > 0$ so that for all multi-indices $|\mathbf{n}| \leq 3$, and $b \in [0, \beta_1]$ we have

$$\cosh(b|\mathbf{y}|)D^{\mathbf{n}}Q(\mathbf{y}) \in L^\infty.$$

Equation (1) is invariant under spatial translations and rotations. So if we let $\mathcal{R}[\theta](x_1, x_2) := (\cos(\theta)x_1 - \sin(\theta)x_2, \sin(\theta)x_1 + \cos(\theta)x_2)$ and if $\mathbf{v}(\theta) := \mathcal{R}[\theta]\mathbf{i}$ then

$$Q(\mathcal{R}[-\theta](\mathbf{x} - ct\mathbf{v}(\theta) - \mathbf{r}_0))$$

is another pulse solution which moves in the direction $\mathbf{v}(\theta)$, *i.e.* at angle θ .

If we insert

$$u(\mathbf{x}, t) = Q(\mathcal{R}[-\theta](\mathbf{x} - ct\mathbf{v}(\theta) - \mathbf{r}_0)) + W(\mathcal{R}[-\theta](\mathbf{x} - ct\mathbf{v}(\theta) - \mathbf{r}_0), t)$$

into (1) we find

$$W_t = AW + O(|W|^2)$$

where the linearization of (1) about the pulse is:

$$A := c\mathbf{i} \cdot \nabla + \mathcal{D}\Delta + F'(Q(\mathbf{y})).$$

We use \mathbf{y} to denote the independent spatial variable of W , which corresponds to the frame moving with the pulse. Note that the kernel of A contains Q_{y_1} and Q_{y_2} due to the translation invariance of (1) and also $\mathbf{y}^\perp \cdot \nabla Q$ because of its rotational invariance. (Here $(y_1, y_2)^\perp := (-y_2, y_1)$.) We assume that these fully characterize the kernel and that the rest of the spectrum is stable. To wit:

Hypothesis 2. *The spectrum of A viewed as an unbounded operator on L^5 , consists of a triple eigenvalue at zero due to the translation and rotation invariance of the problem and the rest which lies in the set*

$$S := \{\lambda \mid \Re\lambda < -\alpha, \quad |\arg(\lambda)| > \pi - \varphi_0\}$$

where $\alpha > 0$ and $\varphi_0 \in (0, \pi/2)$. The center eigenspace is

$$E^c := \text{span} \{ \nabla Q(\mathbf{y}), \mathbf{y}^\perp \cdot \nabla Q(\mathbf{y}) \}$$

with associated spectral projection Π^c . The strong stable eigenspace is denoted $E^s := \ker \Pi^c$ with projections $\Pi^s := 1 - \Pi^c$. Moreover the spectral projection onto E^c is given by $\Pi^c f = \langle \psi^\dagger, f \rangle$ where the adjoint eigenfunction ψ^\dagger has $\cosh(b|\mathbf{x}|)\psi^\dagger(\mathbf{x}) \in W^{1,\infty}$ for $b \in [0, \beta_2]$, for some $\beta_2 > 0$. Note that $\langle \cdot, \cdot \rangle$ denotes the L^2 inner product.

This hypothesis has two important consequences. First, since A is elliptic, this implies (see [2, 16]):

Lemma 3. *There is a positive constant M such that the resolvent estimate*

$$(1 + |\lambda|) \|(\lambda - A)^{-1}\|_{L^5 \rightarrow L^5} \leq M$$

is true for all $\lambda \notin S$. Additionally A generates an analytic semigroup e^{At} on L^5 .

Second, the conjugated operator $A_b := \cosh(b|\mathbf{x}|)A[\operatorname{sech}(b|\mathbf{x}|)\cdot]$ is a small bounded perturbation of A when b is not too large. The spectral properties of A_b as an unbounded operator on L^5 coincide with those of A on

$$L_b^5 := \{f(\mathbf{y}) : \cosh(b|\mathbf{y}|)f(\mathbf{y}) \in L^5\}.$$

Since the center eigenfunctions of A are, by hypothesis, in L_b^5 , we know that the center eigenspace of A_b is the same as that of A . Thus we have:

Lemma 4. *There exists $\beta_3 > 0$ so that Hypothesis 2 and Lemma 3 remain true when L^5 is replaced by the exponentially weighted space L_b^5 for $b \in [0, \beta_3]$. All estimates are uniform in b .*

In what follows, let

$$\beta := \min\{\beta_1, \beta_2, \beta_3\} \quad \text{and} \quad b_0 = \beta/2.$$

Hypothesis 2, Lemma 3 and Lemma 4 can be used to show, for instance, that the pulse Q is stable with asymptotic phase for small initial perturbations in

$$W_b^{8/5,5} := \left\{f(\mathbf{y}) : \cosh(b|\mathbf{y}|)f(\mathbf{y}) \in W^{8/5,5}\right\}$$

and $b \in [0, b_0]$ (see [13]).

1.2 Description of main results

We are interested, however, in multipulse solutions. In the planar setting, pulses can approach or separate from another obliquely and this complicates our analysis in comparison with our work in one dimensional problems in [19, 21]. For instance, two pulses may be separated by a great distance while at the same time be moving along nearly parallel trajectories; the effect of each pulse on the other is small, but acts over very long times. Thus we must carefully quantify the interaction between pulses. To this end, we define for $j = 1, 2$

$$\mathbf{r}_j^*(t) := \mathbf{r}_j^*(t; \theta_{j0}, \mathbf{r}_{j0}) := \mathbf{r}_{j0} + ct\mathbf{v}(\theta_{j0})$$

which gives the location of a pulse indexed by j if it travels along a straight line pointing in the direction θ_{j0} and is initially located at \mathbf{r}_{j0} . Let $\Delta\mathbf{r}_0 := \mathbf{r}_{20} - \mathbf{r}_{10}$, $\Delta\mathbf{v}_0 := \mathbf{v}(\theta_{20}) - \mathbf{v}(\theta_{10})$,

$$\mu^*(t) := \mu^*(t; \theta_{10}, \theta_{20}, \mathbf{r}_{10}, \mathbf{r}_{20}) = |\mathbf{r}_2^*(t) - \mathbf{r}_1^*(t)|$$

and

$$\mu_0^* := \min_{t \geq 0} \mu^*(t).$$

Elementary considerations show that this minimum is achieved at $T^* := \max\left\{\frac{-\Delta\mathbf{r}_0 \cdot \Delta\mathbf{v}_0}{c|\Delta\mathbf{v}_0|^2}, 0\right\}$ and if $T^* > 0$ then

$$\mu_0^* = \left[|\Delta\mathbf{r}_0|^2 - \left(\frac{\Delta\mathbf{r}_0 \cdot \Delta\mathbf{v}_0}{|\Delta\mathbf{v}_0|}\right)^2\right]^{1/2}.$$

Otherwise $\mu_0^* = |\Delta\mathbf{r}_0|$. Likewise, for all $t \geq 0$, we have

$$\mu^*(t) \geq k^*t$$

where

$$k^* := c|\Delta\mathbf{v}_0|\frac{\mu_0^*}{|\Delta\mathbf{r}_0|}.$$

The constant k^* is optimal.

We define the first set of function spaces on which we will work. For $0 < T \leq \infty$, $a \geq 0$ and $b \geq 0$:

$$\begin{aligned} X_{a,b}[T] &:= \{f(\mathbf{x}, t) : e^{at} \cosh(b|\mathbf{x}|)f(\mathbf{x}, t) \in L^5([0, T]; L^5(\mathbf{R}^2))\} \\ X'_{a,b}[T] &:= \{f(\mathbf{x}, t) : e^{at} \cosh(b|\mathbf{x}|)f(\mathbf{x}, t) \in L^5([0, T]; W^{2,5}(\mathbf{R}^2)) \cap W^{1,5}([0, T]; L^5(\mathbf{R}^2))\}. \end{aligned}$$

See Section 2 for further discussion of these particular choices.

Our first result states that states that if the initial data for (1) is sufficiently close to the linear superposition of two pulses which, in the absence of any interaction, would not pass too close to one another, then the solution is of (1) is asymptotically close to the linear superposition of two (possibly different) pulses. We refer to this situation as an *exit*. Precisely:

Theorem 5. *Given Hypothesis 1 and 2, for all $\epsilon_0 > 0$ there exist positive constants δ_{exit} , M_0 and \mathcal{K}_0 so that*

$$\mu_0^* \geq M_0, \quad |\Delta\mathbf{v}_0| \geq \epsilon_0$$

and

$$\left\| u_0 - \sum_{j=1}^2 Q(\mathcal{R}[-\theta_{j0}] (\cdot - \mathbf{r}_{j0})) \right\|_{W^{8/5,5}} \leq \delta_{exit}$$

imply that $u(\mathbf{x}, t)$, the solution of (1) with initial data u_0 , satisfies

$$\left\| u(\cdot, \diamond) - \sum_{j=1}^2 Q(\mathcal{R}[-\theta_j(\diamond)] (\cdot - \mathbf{r}_j(\diamond))) \right\|_{X'_{a,0}[\infty]} \leq \mathcal{K}_0$$

where $a = \min\{\alpha/2, b_0 k^*/8\}$. The functions $\mathbf{r}_j(t)$ and $\theta_j(t)$ ($j = 1, 2$) are defined by

$$\theta_j(t) = \theta_{j0} + \phi_j(t)$$

and

$$\mathbf{r}_j(t) = \mathbf{r}_{j0} + \mathbf{p}_j(t) + \int_0^t \mathcal{R}[\theta_j(s)] \mathbf{c} ds$$

where \mathbf{p}_j and θ_j are C^1 functions¹ for which

$$\|\mathbf{p}_j(\diamond)\|_{C^1(\mathbf{R}^+)} + \|\phi(\diamond)\|_{C^1(\mathbf{R}^+)} \leq \mathcal{K}_0.$$

Remark 6. *In the proof, we find that $M_0 = -c_1 \ln(|\Delta\mathbf{v}_0|/2) + c_2$ for constants $c_1, c_2 > 0$.*

Our second main result concerns pulses which *shoot* in towards one another from spatial infinity. We work with the function spaces

$$\begin{aligned} Z_{\eta,b} &:= \{f(\mathbf{x}, t) : e^{\eta t} \cosh(b|\mathbf{x}|)f(\mathbf{x}, t) \in L^5([-\infty, 0]; L^5(\mathbf{R}^2))\} \\ Z'_{\eta,b} &:= \{f(\mathbf{x}, t) : e^{\eta t} \cosh(b|\mathbf{x}|)f(\mathbf{x}, t) \in L^5([-\infty, 0]; W^{2,5}(\mathbf{R}^2)) \cap W^{1,5}([-\infty, 0]; L^5(\mathbf{R}^2))\}. \end{aligned}$$

Our next theorem states that for suitable choices of $\mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}$, and θ_{20} that there is a solution, defined for all $t \leq 0$, which is exponentially close to the sum of two pulses moving along trajectories \mathbf{r}_1^* and \mathbf{r}_2^* :

¹Note that we use the supremum norm for $C^1[0, T]$.

Theorem 7. *Given Hypothesis 1 and 2, for all $\epsilon_0 > 0$ there exist positive constants M_1 and \mathcal{K}_1 so that if*

$$|\Delta \mathbf{v}_0| \geq \epsilon_0, \quad \Delta \mathbf{r}_0 \cdot \Delta \mathbf{v}_0 \leq 0 \quad \text{and} \quad |\Delta \mathbf{r}_0| \geq M_1$$

then there exists $\phi(\mathbf{x}, t) = \phi(\mathbf{x}; \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20})$ which solves (1) and satisfies:

$$\left\| \phi(\cdot, \diamond) - \sum_{j=1}^2 Q(\mathcal{R}[-\theta_{j0}] (\cdot - \mathbf{r}_j^*(\diamond))) \right\|_{Z'_{\eta,0}} \leq \mathcal{K}_1.$$

Here $\eta = -b_0 c |\Delta \mathbf{v}_0| / 8$.

Moreover, ϕ has a decomposition into spatially localized pieces. That is, there are $W_1^*, W_2^* \in Z'_{\eta, b_0}$ (with norms less than \mathcal{K}_1) so that

$$\phi(\mathbf{x}, t) = \sum_{j=1}^2 Q(\mathcal{R}[-\theta_{j0}] (\mathbf{x} - \mathbf{r}_j^*(t))) + W_j^*(\mathcal{R}[-\theta_{j0}] (\mathbf{x} - \mathbf{r}_j^*(t)), t).$$

W_1^* and W_2^* (and therefore ϕ) are differentiable in their dependence on the parameters $\mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20}$.

Remark 8. *The condition $\Delta \mathbf{r}_0 \cdot \Delta \mathbf{v}_0 \leq 0$ just implies that for $t \leq 0$, the pulses are closest at $t = 0$, not before.*

Our final result concerns the stability of the shooting solutions.

Theorem 9. *There exist positive constants δ_{shoot}, T_3 and \mathcal{K}_2 so that if $T \geq T_3$ and*

$$\|u_0 - \phi(\cdot, -T; \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20})\|_{W^{8/5,5}} = \delta \leq \delta_{shoot}$$

then $u(\mathbf{x}, t)$, the solution of (1) with initial data u_0 , satisfies

$$\left\| u(\cdot, \diamond) - \phi(\cdot, \diamond - T; \tilde{\mathbf{r}}_{10}, \tilde{\mathbf{r}}_{20}, \tilde{\theta}_{10}, \tilde{\theta}_{20}) \right\|_{X'_{\alpha/2,0}[T-T_3]} \leq \mathcal{K}_2 \delta$$

where

$$\sum_j |\mathbf{r}_{j0} - \tilde{\mathbf{r}}_{j0}| + |\theta_{j0} - \tilde{\theta}_{j0}| \leq \mathcal{K}_2 \delta.$$

Note that this theorem implies:

Corollary 10. *The function $\phi(\mathbf{x}, t; \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20})$ is the unique function with properties described in Theorem 7.*

1.3 Interaction manifolds

Given the freedom in choosing $\mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}$ and θ_{20} in our main results we define the following *interaction manifolds* (defined for fixed ϵ_0):

$$\mathcal{M}_{exit} := \{Q(\mathcal{R}[\theta_{10}](\cdot - \mathbf{r}_{10})) + Q(\mathcal{R}[\theta_{20}](\cdot - \mathbf{r}_{20})) : \Delta \mathbf{v}_0 \geq \epsilon_0 \quad \text{and} \quad \mu_0^* \geq M_0\}$$

$$\mathcal{M}_{shoot} := \{\phi(\cdot, 0; \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20}) : \Delta \mathbf{v}_0 \geq \epsilon_0, \quad \Delta \mathbf{r}_0 \cdot \Delta \mathbf{v}_0 \leq 0 \quad \text{and} \quad |\Delta \mathbf{r}_0| \geq M_1\}.$$

Since Q and ϕ are in $W^{8/5,5}$ and depend differentiably on $\mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}$ and θ_{20} , these are six-dimensional C^1 manifolds in $W^{8/5,5}$. Each has a boundary. Note that \mathcal{M}_{shoot} is an invariant manifold for (1) since ϕ solves this equation exactly, whereas \mathcal{M}_{exit} is not. Nevertheless, our main results indicate that both are attracting sets for (1). That is we have the following corollaries of our main theorems (and also (6)):

Corollary 11. *Suppose that*

$$\text{dist}_{W^{8/5,5}}(u(x, 0), \mathcal{M}_{\text{exit}}) \leq \delta_{\text{exit}}.$$

Then there exists a $a > 0$ so that

$$\text{dist}_{W^{8/5,5}}(u(x, t), \mathcal{M}_{\text{exit}}) \leq \mathcal{K}_0 e^{-at}.$$

for all $t \geq 0$.

Corollary 12. *Suppose that*

$$\text{dist}_{W^{8/5,5}}(u(x, 0), \mathcal{M}_{\text{shoot}}) = \delta \leq \delta_{\text{shoot}}.$$

Then

$$\text{dist}_{W^{8/5,5}}(u(x, t), \mathcal{M}_{\text{shoot}}) \leq \mathcal{K}_2 \delta e^{-\frac{b_0}{8} c |\Delta \mathbf{v}_0| t}.$$

for all $0 \leq t \leq T$. Here $T > 0$ is a constant which depends only upon the parameters \mathbf{r}_{10} , \mathbf{r}_{20} , θ_{10} and θ_{20} which minimize $\text{dist}_{W^{8/5,5}}(u(x, 0), \mathcal{M}_{\text{shoot}})$.

We make two final remarks, the first of which is paraphrased from [19].

Remark 13. *Corollary 12 implies that collisions between two pulses are well-defined scattering problems. By this, we mean that strong interactions which take place during collisions will not be affected by specific choices for the initial data which lead to this collision. Moreover Corollary 11 tells us that if there is an interaction which results after a finite amount of time in two pulses which are moving away from one another, that this state will persist for all time. That is to say, these results indicate that to study strong collisions one only needs finite time results.*

Remark 14. *Theorem 7 implies that if we take $\Delta \mathbf{r}_0$ to infinity while keeping $\Delta \mathbf{v}_0$ fixed, that*

$$\|\phi(\cdot, 0; \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20}) - Q(\mathcal{R}[\theta_{10}](\cdot - \mathbf{r}_{10})) - Q(\mathcal{R}[\theta_{20}](\cdot - \mathbf{r}_{20}))\|_{W^{8/5,5}} \rightarrow 0$$

which is to say that

$$\text{dist}(\mathcal{M}_{\text{shoot}}, \mathcal{M}_{\text{exit}}) = 0.$$

This implies that there are choices for \mathbf{r}_{10} , \mathbf{r}_{20} , θ_{10} , and θ_{20} so that $\phi(\cdot, t; \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20})$ is in the attracting region for $\mathcal{M}_{\text{exit}}$ for some $t \leq 0$. Thus there are global in time solutions of (1) which converge exponentially quickly for $|t| \rightarrow \infty$ to the linear superposition of two pulses.

1.4 The embedding and general strategy

Our strategy is to embed (1) into a larger system within which the multipulse problem can be viewed, in some sense, as a small perturbation of single pulse problem. To wit, we consider

$$\begin{aligned} U_t &= \mathcal{D}\Delta U + F(U) + G_1(U, V) \\ V_t &= \mathcal{D}\Delta V + F(V) + G_2(U, V) \end{aligned} \tag{2}$$

where we choose G_1 and G_2 so that

$$F(U) + F(V) + G_1(U, V) + G_2(U, V) = F(U + V).$$

With this, if U and V solve (2), then

$$u = U + V$$

solves (1). Note that this idea is very similar to the ‘‘freezing’’ method used in [4]. The idea is then to show that there are solutions of (2) roughly of the form

$$\begin{aligned} U &= Q(\mathcal{R}[-\theta_{10}](\mathbf{x} - \mathbf{r}_1^*(t))) + \text{‘‘small’’} \\ V &= Q(\mathcal{R}[-\theta_{20}](\mathbf{x} - \mathbf{r}_2^*(t))) + \text{‘‘small’’} \end{aligned}$$

where the parameters are θ_{j0} and \mathbf{r}_{j0} are taken so that μ_0^* is large.

We select G_1 and G_2 as follows. Suppose that at time t our pulses are “located” at $\mathbf{r}_1(t)$ and $\mathbf{r}_2(t)$. Let $L(t)$ be the perpendicular bisector of the segment connecting these two points. The line $L(t)$ divides \mathbf{R}^2 into a disjoint union of two sets $\Sigma_1(t)$ (which contains the pulse labeled “1”) and $\Sigma_2(t)$ (which contains the other). For concreteness, we assume $L(t) \subset \Sigma_1(t)$.

Let $\chi_j(\mathbf{x}; \mathbf{r}_1(t), \mathbf{r}_2(t))$, $j = 1, 2$ be a C^∞ partition of unity subordinate to the sets $\Sigma_j(t)$ and which have derivatives whose support lies in $\{\mathbf{x} : \text{dist}(\mathbf{x}, L(t)) \leq 1\}$. More compactly we will write these as $\chi_1(\mathbf{x}, t)$ and $\chi_2(\mathbf{x}, t)$. We set:

$$G(U, V) := (F(U + V) - F(U) - F(V))$$

and

$$G_j(U, V) = \chi_j(\mathbf{x}, t)G(U, V).$$

We are thinking of U as being the pulse which lies in Σ_1 and V as being the one in Σ_2 .

Our choice for G is motivated by the following heuristic. A straightforward application of Taylor’s theorem, together with the fact that $F(0) = 0$ implies:

$$|G(U, V)| \leq C|U||V|.$$

Now suppose that $U = Q_1 + W_1$ and $V = Q_2 + W_2$ where Q_1 and Q_2 are pulses and W_1 and W_2 are error functions. Thus

$$|G_1(U, V)| \leq C\chi_1|Q_1||Q_2| + C\chi_1|Q_1||W_2| + C\chi_1|Q_2||W_1| + C\chi_1|W_1||W_2|.$$

Of the four terms here, three are “small”.

- Q_1 and Q_2 are exponentially localized and so $|Q_1||Q_2|$ is exponentially small in the separation distance.
- The term $|W_1||W_2|$ is quadratic and thus very small if the error functions are small.
- The term $\chi_1|Q_2|$ is clearly bounded by the maximum value of Q_2 evaluated on the region Σ_1 , which is to say that Q_2 is evaluated far from its center. Thus $\chi_1|Q_2||W_1|$ is small as well.

The remaining term $\chi_1|Q_1||W_2|$ is not small, at least not for general functions W_2 . If W_2 is localized near the pulse Q_2 then we can conclude that this term is exponentially small in the pulse separation distance. Thus if we work with spaces of localized functions, we can handle this term. We wish to allow arbitrary perturbations of the pulses, however, so this restriction is unsatisfying. We are able to circumvent this difficulty by making the observation that even if W_2 is not localized, $\chi_1|Q_1||W_2|$ is. It turns out that this allows us to treat the term as if it were small.

The following toy model linear algebra problem demonstrates the core idea of why this is so. Let \mathcal{A} be an invertible n by n matrix (which is our stand-in here for the linearization A). Suppose we are trying to solve the equation

$$(\mathcal{A} + \mathcal{B})w = j \tag{3}$$

where \mathcal{B} is a matrix that has the following properties:

1. $\mathcal{B} : \mathbf{R}^n \rightarrow X_b$, where X_b is a subspace of \mathbf{R}^n .
2. $\|\mathcal{B}\|_{X_b \rightarrow X_b} = \epsilon \ll 1$.

(The matrix \mathcal{B} is playing the role of the perturbation to A made by $\chi_1|Q_1||W_2|$ and X_b corresponds to the space of “exponentially decay functions.”) If in addition we know that

$\mathcal{A}^{-1} : X_b \rightarrow X_b$, then the fact that \mathcal{B} is small on X_b means that $\mathcal{A} + \mathcal{B}$ is invertible on X_b . We denote this inverse by $(\mathcal{A} + \mathcal{B})_{X_b}^{-1}$. Now consider the augmented system

$$\begin{aligned} \mathcal{A}w^i &= j \\ (\mathcal{A} + \mathcal{B})w^l &= -\mathcal{B}w^i. \end{aligned} \tag{4}$$

The first equation can be solved for any j in \mathbf{R}^n . Since \mathcal{B} maps everything into X_b , we can solve the second equation. Moreover, adding these two equations shows that $w = w^i + w^l$ solves (3). Specifically, the solution map is:

$$\mathcal{G} := \mathcal{A}^{-1} - (\mathcal{A} + \mathcal{B})_{X_b}^{-1} \mathcal{B} \mathcal{A}^{-1}.$$

Thus \mathcal{G} is $(\mathcal{A} + \mathcal{B})^{-1}$ on all of \mathbf{R}^n .

The remainder of this paper is organized as follows. In Section 2 we discuss the functional setting for our results. We compute the ‘‘almost’’ linearization of (2) about a multipulse solution in Section 3 and collect a number of useful estimates. Section 4 concerns exit solutions and in particular contains the proof of Theorem 5. Section 5 is about the shooting manifold and has the proofs of Theorems 7 and 9 and Corollary 10. In the Appendix we prove a local in time existence theorem for solutions of a system needed in Section 4.

Acknowledgements: The author would like to thank Thierry Gallay for suggesting the choice of collective coordinates used in Section 4. Additional gratitude is due to the Drexel University PDE group and Arnd Scheel. Finally, the NSF generously funded this work under grant DMS 0807738.

2 Functional setting and maximal regularity

In this section we discuss important properties of A and solutions of equations of the form

$$W_t = AW + J(t) \quad \text{and} \quad W(t=0) = W_0. \tag{5}$$

Specifically, we summarize the maximal regularity results for Sobolev spaces, found in Section III.4 of [2]. The domain of A viewed as an operator on L^p is of course $W^{2,p}$ and so we set

$$Y := L^p([0, T]; L^p) \quad \text{and} \quad Y' := W^{1,p}([0, T]; L^p(\mathbf{R}^2)) \cap L^p([0, T]; W^{2,p}(\mathbf{R}^2)).$$

Hypothesis 2 and Theorem 4.10.7 in [2] allow us to conclude that there is a continuous solution map Γ (given explicitly by the Duhamel formula)

$$\Gamma : Y \times E_{1-1/p,p} \rightarrow Y'.$$

That is, if $W = \Gamma(J, W_0)$ then W satisfies (5). The space

$$E_{1-1/p,p} = (L^p(\mathbf{R}^2), W^{2,p}(\mathbf{R}^2))_{1-1/p,p},$$

where $(\cdot, \cdot)_{\theta,q}$ is the real interpolation functor. We have (see, for instance, [20]) that

$$(L^p(\mathbf{R}^2), W^{2,p}(\mathbf{R}^2))_{1-1/p,p} = W^{2-2/p}(\mathbf{R}^2).$$

Also, Theorem 4.10.2 in [2] tells us

$$Y' \subset\subset BUC([0, T]; W^{2-2/p}(\mathbf{R}^2)).$$

Our equation is nonlinear, and so at the very least, we would like $W^{2-2/p} \subset\subset C(\mathbf{R}^2)$; this together with the above inclusion implies that Y' is an algebra. At several points, we require $W^{2-2/p} \subset\subset C^1(\mathbf{R}^2)$. From Sobolev embedding we get this inclusion when $1 - 2/p > 2/p$ or rather $p > 4$. We take $p = 5$.

Now we restate Theorem 4.10.7 and Remark 4.10.9 (a) here, adapted to our problem:

Theorem 15. *There exists $C > 0$ so that for all $b \in [0, b_0]$ the following are true:*

1. *For all $a \in [0, \alpha/2]$ and $T \in (0, \infty]$, the map*

$$\Gamma_1(W^s, J) := e^{At}\Pi^s W^s + \int_0^t e^{A(t-\tau)}\Pi^s J(\tau)d\tau$$

is bounded from $(E^s \cap W_b^{8/5,5}) \times X_{a,b}[T]$ to $X'_{a,b}[T]$. Additionally, it satisfies

$$\|\Gamma_1(W^s, J)\|_{X'_{a,b}[T]} \leq C \left(\|W^s\|_{W_b^{8/5,5}} + \|J\|_{X_{a,b}[T]} \right)$$

and $W(\mathbf{y}, t) := \Gamma_1(W^s, J)$ solves (for $t \geq 0$ a.e.)

$$W_t = AW + \Pi^s J, \quad W(\mathbf{y}, 0) = W^s \in E^s.$$

2. *For all $\eta < 0$, the map*

$$\Gamma_2(J) := \int_{-\infty}^t e^{A(t-\tau)}J(\tau)d\tau$$

is bounded from $Z_{\eta,b}$ to $Z'_{\eta,b}$. Additionally, it satisfies

$$\|\Gamma_2(J)\|_{Z'_{\eta,b}} \leq C \|J\|_{Z_{\eta,b}}$$

and $W(\mathbf{y}, t) := \Gamma_2(J)$ solves (for $t \leq 0$ a.e.)

$$W_t = AW + J.$$

3. *For all $a \in [0, \alpha/2]$ and $T \in (0, \infty]$, the map*

$$\Gamma_3(W^s, J) := e^{At}\Pi^s W^s + \int_0^t e^{A(t-\tau)}\Pi^s J(\tau)d\tau - \int_t^T e^{A(t-\tau)}\Pi^c J(\tau)d\tau$$

is bounded from $(E^s \cap W_b^{8/5,5}) \times X_{a,b}[T]$ to $X'_{a,b}[T]$. Additionally, it satisfies

$$\|\Gamma(W^s, J)\|_{X'_{a,b}[T]} \leq C \left(\|W^s\|_{W_b^{8/5,5}} + \|J\|_{X_{a,b}[T]} \right)$$

and $W(\mathbf{y}, t) := \Gamma(W^s, J)$ solves (for $t \geq 0$ a.e.)

$$W_t = AW + \Pi^s J, \quad \Pi^s W(\mathbf{y}, 0) = W^s \in E^s \quad \text{and} \quad \Pi^c W(\mathbf{y}, T) = 0.$$

Remark 16. *The first of these will be used to prove the stability of exiting pulses, the second the existence of shooting solutions and the last the stability of shooting solutions. For a further discussion of these results and how they follow from [2], see [19].*

Finally, we note that Theorem 4.10.2 gives:

$$\begin{aligned} X'_{a,b}[T] &\subset\subset \left\{ f(\mathbf{x}, t) : e^{at}f(\mathbf{x}, t) \in BUC([0, T]; W_b^{8/5,5}) \right\} \\ Z'_{\eta,b} &\subset\subset \left\{ f(\mathbf{x}, t) : e^{\eta t}f(\mathbf{x}, t) \in BUC([-\infty, 0]; W_b^{8/5,5}) \right\}. \end{aligned} \tag{6}$$

3 The “almost” linearization of (2) around a multi-pulse

3.1 Reduction

We consider (2) with initial conditions

$$U(\mathbf{x}, 0) = Q(\mathcal{R}[-\theta_{10}](\mathbf{x} - \mathbf{r}_{10})) + W_{10}(\mathcal{R}[-\theta_{10}](\mathbf{x} - \mathbf{r}_{10}))$$

$$V(\mathbf{x}, 0) = Q(\mathcal{R}[-\theta_{20}](\mathbf{x} - \mathbf{r}_{20})) + W_{20}(\mathcal{R}[-\theta_{20}](\mathbf{x} - \mathbf{r}_{20})).$$

The following lemma tells us that we can without loss of generality assume that W_{10} and W_{20} lie in E^s .

Lemma 17. *There is a smooth map $\mathcal{I} = (\mathcal{I}_1, \mathcal{I}_2, \mathcal{I}_3) : W^{8/5,5} \longrightarrow \mathbf{R}^2 \times \mathbf{R} \times E^s$ with the property that*

$$\mathcal{I}(0) = (0, 0, 0)$$

and if $\|W\|_{W^{8/5,5}}$ is sufficiently small, then

$$Q(\mathbf{x}) + W(\mathbf{x}) = Q(\mathcal{R}[\mathcal{I}_2(W)](\mathbf{x} - \mathcal{I}_1(W))) + \mathcal{I}_3(W)(\mathcal{R}[\mathcal{I}_2(W)](\mathbf{x} - \mathcal{I}_1(W))).$$

Proof. The proof follows immediately from applying the implicit function theorem to the map:

$$\Theta(\mathcal{I}, W) = Q(\mathbf{x}) + W(\mathbf{x}) - Q(\mathcal{R}[\mathcal{I}_2](\mathbf{x} - \mathcal{I}_1)) - \mathcal{I}_3(\mathcal{R}[\mathcal{I}_2](\mathbf{x} - \mathcal{I}_1)).$$

□

Now we let

$$\begin{aligned} U(\mathbf{x}, t) &= Q(\mathbf{z}_1) + W_1(\mathbf{y}_1, t) \\ V(\mathbf{x}, t) &= Q(\mathbf{z}_2) + W_2(\mathbf{y}_2, t) \end{aligned} \tag{7}$$

where

$$\begin{aligned} \mathbf{y}_1 &= \mathcal{R}[-\theta_{10}](\mathbf{x} - \mathbf{r}_1(t)) & \text{and} & & \mathbf{z}_1 &= \mathcal{R}[-\theta_1(t)](\mathbf{x} - \mathbf{r}_1(t)) \\ \mathbf{y}_2 &= \mathcal{R}[-\theta_{20}](\mathbf{x} - \mathbf{r}_2(t)) & \text{and} & & \mathbf{z}_2 &= \mathcal{R}[-\theta_2(t)](\mathbf{x} - \mathbf{r}_2(t)). \end{aligned}$$

The collective coordinates $\mathbf{r}_j(t)$ and $\theta_j(t)$ are specified in Sections 4 and 5. Note here that the “spatial coordinates” of the pulses Q and the error functions W_1 and W_2 are not all the same. The coordinates on the RHS for U are “centered” at the point $\mathbf{r}_1(t)$, and those for V at $\mathbf{r}_2(t)$. Moreover, the coordinates for the W s have their rotation matrices fixed at their initial values, $\theta_j(0) = \theta_{j0}$.

We use

$$\mathbf{m} := (\mathbf{p}_1, \phi_1, \mathbf{p}_2, \phi_2)$$

(with \mathbf{p}_j, ϕ_j defined as in Theorem 5) to refer to the collective coordinates in the aggregate and define

$$\mu(t) := |\mathbf{r}_2(t) - \mathbf{r}_1(t)|$$

the separation distance of the two pulses.

Inserting (7) into (2), we find:

$$\begin{aligned} \partial_t W_1 &= A W_1 + B_1(\mathbf{m}) W_2 \\ &\quad + \{\mathcal{R}[-\theta_1] \dot{\mathbf{r}}_1 - \dot{\mathbf{c}}_1\} \cdot \nabla Q + \dot{\theta}_1 \mathbf{y}^\perp \cdot \nabla Q + J_1 \end{aligned} \tag{8}$$

$$\begin{aligned} \partial_t W_2 &= A W_2 + B_2(\mathbf{m}) W_1 \\ &\quad + \{\mathcal{R}[-\theta_2] \dot{\mathbf{r}}_2 - \dot{\mathbf{c}}_2\} \cdot \nabla Q + \dot{\theta}_2 \mathbf{y}^\perp \cdot \nabla Q + J_2 \end{aligned} \tag{9}$$

where

$$B_1(\mathbf{m})W_2 := \chi_1(\mathbf{x}, t) [F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))] W_2(\mathbf{y}_2, t)$$

and

$$B_2(\mathbf{m})W_1 := \chi_2(\mathbf{x}, t) [F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_1))] W_1(\mathbf{y}_1, t).$$

These terms correspond to the problematic term $\chi_1 |Q_1|$ from the Introduction.

The term J_1 term is given by

$$J_1 := K_1 + J_{1,mod} + J_{1,int} + J_{1,nl}$$

where

$$\begin{aligned} K_1 := & \left\{ (\mathcal{R}[-\theta_1]\mathbf{r}_1 - \mathbf{c}\mathbf{i}) + \dot{\theta}_1 \mathcal{R}[-\theta_1 + \theta_{10}] \mathbf{y}^\perp \right\} \cdot (\nabla Q(\mathbf{z}_1) - \nabla Q(\mathbf{y})) \\ & + \dot{\theta}_1 \left\{ \mathcal{R}[-\theta_1 + \theta_{10}] \mathbf{y}^\perp - \mathbf{y}^\perp \right\} \cdot \nabla Q(\mathbf{y}) \\ & + \mathcal{R}[\theta_1 - \theta_{10}] \left\{ \mathcal{R}[-\theta_1]\mathbf{r}_1 - \mathbf{c}\mathbf{i} \right\} \cdot \nabla W_1 \end{aligned}$$

$$\begin{aligned} J_{1,mod} := & (\mathcal{R}[\theta_1 - \theta_{10}]\mathbf{c}\mathbf{i} - \mathbf{c}\mathbf{i}) \cdot \nabla W_1 \\ & + \{F'(Q(\mathbf{z}_1)) - F'(Q(\mathbf{y}))\} W_1 \end{aligned}$$

$$\begin{aligned} J_{1,int} := & \chi_1(\mathbf{x}, t) [F(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F(Q(\mathbf{z}_1)) - F(Q(\mathbf{z}_2))] \\ & + \chi_1(\mathbf{x}, t) [(F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_1))) W_1] \end{aligned}$$

$$\begin{aligned} J_{1,nl} := & F(Q(\mathbf{z}_1) + W_1) - (F(Q(\mathbf{z}_1)) + F'(Q(\mathbf{z}_1))W_1) \\ & + \chi_1(\mathbf{x}, t) N_{ct}(W_1, W_2(\mathbf{y}_2, t)) \end{aligned}$$

and

$$\begin{aligned} & N_{ct}(W_1, W_2) \\ := & F(Q(\mathbf{z}_1) + Q(\mathbf{z}_2) + W_1 + W_2) - F(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2))(W_1 + W_2) \\ & - (F(Q(\mathbf{z}_1) + W_1) - F(Q(\mathbf{z}_1)) - F'(Q(\mathbf{z}_1))W_1) \\ & - (F(Q(\mathbf{z}_2) + W_2) - F(Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))W_2). \end{aligned}$$

The terms in K_1 and $J_{1,mod}$ are small due to the choice of the collective coordinates, those in $J_{1,int}$ due to the types interactions described in the introduction and those in $J_{1,nl}$ because they are nonlinear. We define J_2 analogously.

Remark 18. *In the definitions of B_j and J_j we use \mathbf{y} as the independent spatial variable for both W_1 and W_2 . If we do not specify the spatial coordinate of a function, it is assumed to be \mathbf{y} . When we have \mathbf{y}_2 (or \mathbf{x} , \mathbf{z}_1 , \mathbf{z}_2) in the equation for W_1 , it is implicit that we are viewing \mathbf{y}_2 as a function of \mathbf{y} and t . For instance when we write $W_2(\mathbf{y}_2, t)$ in $B_1(\mathbf{m})W_2$ we have*

$$\mathbf{y}_2 = \mathbf{y}_2(\mathbf{y}, t) = \mathcal{R}[\theta_{10} - \theta_{20}]\mathbf{y} + \mathcal{R}[-\theta_{20}] (\mathbf{r}_2(t) - \mathbf{r}_1(t)).$$

In particular, we point out the B_j are nonlocal operators.

3.2 Estimates on B_j and J_j

The following lemma, though easy to prove, is key to our strategy:

Lemma 19. For $j = 1, 2$ and for $0 \leq b' \leq b \leq b_0$ we have

$$\| [B_j(\mathbf{m}(t))V](\cdot) \|_{L_b^5} \leq C e^{-\frac{b'}{2}\mu(t)} \| V(\cdot) \|_{L_{b'}^5}.$$

The constant $C > 0$ is independent of b' and of \mathbf{m} .

Proof. We carry out the details for $j = 1$. First notice that since F is C^2 and Q is exponentially localized we have:

$$\begin{aligned} |B_1(\mathbf{m})| &= \chi_1(\mathbf{x}, t) |F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))| \\ &\leq C \chi_1(\mathbf{x}, t) |Q(\mathbf{z}_1)| \\ &\leq C \chi_1(\mathbf{x}, t) e^{-b_0|\mathbf{z}_1|} \\ &\leq C \chi_1(\mathbf{x}, t) e^{-b_0|\mathbf{y}|}. \end{aligned}$$

Therefore

$$e^{b|\mathbf{y}|} |[B_j V](\mathbf{y})| \leq C \chi_1(\mathbf{x}, t) |V(\mathbf{y}_2)| = C \chi_1(\mathbf{x}, t) e^{-b'|\mathbf{y}_2|} e^{b'|\mathbf{y}_2|} |V(\mathbf{y}_2)|. \quad (10)$$

Next,

$$\chi_1(\mathbf{x}, t) e^{-b'|\mathbf{y}_2|} \leq \exp\left(-b' \inf_{\mathbf{x} \in \text{supp}\chi_1(t)} |\mathbf{y}_2(t)|\right) \leq \exp\left(-b' \inf_{\mathbf{x} \in \text{supp}\chi_1(t)} |\mathbf{x} - \mathbf{r}_2(t)|\right).$$

By assumption χ_1 is zero outside $\Sigma_1 \cup \{\mathbf{x} : \text{dist}(\mathbf{x}, L(t)) \leq 1\}$ and Σ_1 is the set of points ‘‘on same side’’ of $L(t)$ as \mathbf{r}_1 . Also recall that $L(t)$ is equidistant from \mathbf{r}_1 and \mathbf{r}_2 . Thus we conclude that the minimum value of $|\mathbf{x} - \mathbf{r}_2|$ in Σ_1 occurs halfway between \mathbf{r}_1 and \mathbf{r}_2 , *i.e.* at $\mathbf{x}_0 = (\mathbf{r}_1(t) + \mathbf{r}_2(t))/2$. The minimum value of $|\mathbf{x} - \mathbf{r}_2|$ in the slightly larger set $\text{supp}\chi_1$ must occur within a distance of 1 of \mathbf{x}_0 and so:

$$\chi_1(\mathbf{x}, t) e^{-b'|\mathbf{y}_2|} \leq C e^{-\frac{b'}{2}\mu(t)}.$$

Taking the L^5 norm of (10) and using this last inequality finishes the proof. \square

Notice two important features of this Lemma. First, setting $b' = 0$ and $b = b_0$, we see $B_j V$ decays in space more rapidly than V does, although the norm of B_j as a map from L^5 to $L_{b_0}^5$ is $O(1)$. Second, if $b' = b = b_0$ then the norm of B_j as a map from $L_{b_0}^5$ to $L_{b_0}^5$ is exponentially small in the separation distance.

The next set of lemmas give L^5 estimates for the terms which comprise J_j .

Lemma 20. There exists a constant $C > 0$ so that for $j = 1, 2$, $t \geq 0$ and $|\phi_j(t)| \leq \pi/2$:

$$\| J_{j,mod}(\cdot, t) \|_{L^5} \leq C |\mathbf{m}(t)| \| W_j(\cdot, t) \|_{W^{8/5,5}}.$$

Lemma 21. We have a constant $C > 0$ so that for all $0 \leq b \leq b_0$, $j = 1, 2$ and all $t \geq 0$:

$$\| J_{j,int}(\cdot, t) \|_{L_b^5} \leq C e^{-\frac{b_0}{2}\mu(t)} \left(1 + \| W_1(\cdot, t) \|_{L_b^5}\right).$$

Lemma 22. We have a constant $C > 0$ so that for all $0 \leq b \leq b_0$, $j = 1, 2$ and all $t \geq 0$:

$$\| J_{j,nl}(\cdot, t) \|_{L_b^5} \leq C \left(\| W_1 \|_{W_b^{8/5,5}}^2 + \| W_2 \|_{W_b^{8/5,5}}^2 \right).$$

Lemma 23. There exists $C > 0$ so that for $j = 1, 2$, $t \geq 0$ and $|\phi_j(t)| \leq \pi/2$

$$\| K_j(\cdot, t) \|_{L^5} \leq C |\dot{\mathbf{m}}(t)| (|\mathbf{m}(t)| + \| W_j(\cdot, t) \|_{W^{8/5,5}}).$$

We now prove these in order:

Proof. (Lemma 20) We have $|\mathcal{R}[\theta_1 - \theta_{10}] - id| \leq C|\phi_1|$ and so the first term in $J_{1,mod}$ is handled easily.

For the second term, we use that $F \in C^2$ to see:

$$|F'(Q(\mathbf{z}_1)) - F'(Q(\mathbf{y}))| \leq C|Q(\mathbf{z}_1) - Q(\mathbf{y})|$$

By the mean value theorem there is a \mathbf{y}_* on the line segment connecting \mathbf{z}_1 to \mathbf{y} so that:

$$|Q(\mathbf{z}_1) - Q(\mathbf{y})| = |\mathbf{z}_1 - \mathbf{y}| |\nabla Q(\mathbf{y}_*)| \leq |(\mathcal{R}[\phi_1] - id)| |\mathbf{y}| |\nabla Q(\mathbf{y}_*)| \leq C|\phi_1| |\mathbf{y}| e^{-\beta|\mathbf{y}_*|}.$$

Since $|\mathbf{z}_1| = |\mathbf{y}|$ and \mathbf{y}_* is on the segment connecting them, we have $|\mathbf{y}_*| \geq |\mathbf{y}| \cos(\phi_1/2) \geq \sqrt{2}|\mathbf{y}|/2$. This implies $|\mathbf{y}| e^{-\beta|\mathbf{y}_*|} \leq C$ and we are done. \square

Proof. (Lemma 21) The estimate for the first term goes as follows: We have $F \in C^2$ and $F(0)$ and therefore there exists a constant $C > 0$ so that for all $Q_1, Q_2 \in \mathbf{R}^N$

$$|F(Q_1 + Q_2) - F(Q_1) - F(Q_2)| \leq C|Q_1||Q_2|.$$

The proof can be obtained by the mean value theorem ([21]). Thus

$$\begin{aligned} |\chi_1(\mathbf{x}, t) [F(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F(Q(\mathbf{z}_1)) - F(Q(\mathbf{z}_2))]| \\ \leq C\chi_1(\mathbf{x}, t)|Q(\mathbf{z}_1)||Q(\mathbf{z}_2)| \\ \leq C \left(\sup_{\mathbf{x} \in \text{supp}\chi_1(t)} \exp(-b_0|\mathbf{z}_2|) \right) e^{-\beta|\mathbf{z}_1|} \leq C e^{-\frac{b_0}{2}\mu(t)} e^{-\beta|\mathbf{y}|}. \end{aligned}$$

Note that we have used the same estimate as appeared in the proof of Lemma 19. Since $e^{-\beta|\mathbf{y}|}$ is in L_b^5 we have completed the estimate for the first term.

The proof for the second term is nearly identical to the proof for Lemma 19, though it is easier. To wit, we have

$$\begin{aligned} |\chi_1(\mathbf{x}, t) [(F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_1))) W_1(\mathbf{y})]| \\ \leq C\chi_1(\mathbf{x}, t) |Q(\mathbf{z}_2)| |W_1(\mathbf{y})| \\ \leq C \left(\sup_{\mathbf{x} \in \text{supp}\chi_1(t)} \exp(-b_0|\mathbf{z}_2|) \right) |W_1(\mathbf{y})| \leq C e^{-\frac{b_0}{2}\mu(t)} |W_1(\mathbf{y})|. \end{aligned}$$

\square

Proof. (Lemma 22) This estimate follows from Taylor's theorem applied to F combined with the fact that $W^{8/5,5}$ is an algebra. \square

Proof. (Lemma 23) First:

$$\mathbf{p}_1 = \mathbf{r}_1 - \mathcal{R}[\theta_1(t)]\mathbf{c}\mathbf{i} = \mathcal{R}[\theta_1(t)] (\mathcal{R}[-\theta_1(t)]\mathbf{r}_1(t) - \mathbf{c}\mathbf{i})$$

and

$$\dot{\phi}_1 = \dot{\theta}_1.$$

Then an argument nearly identical to that used in the proof of Lemma 20 shows that for $|\phi_1| \leq \pi/2$ we have

$$|(1 + \mathbf{y}^\perp \cdot) (\nabla Q(\mathbf{z}_1) - \nabla Q(\mathbf{y}))| \leq C|\phi_1(t)|(1 + |\mathbf{y}|)e^{-h|\mathbf{y}|}$$

for a constant $h > 0$. Since $|\mathbf{y}|e^{-h|\mathbf{y}|} \in L^5$ we can estimate the first line in K_1 by $C|\dot{\mathbf{m}}(t)||\mathbf{m}(t)|$.

The estimate for the second line follows from observing that $|\mathbf{y}||Q(\mathbf{y})| \in L^5$ and $|\mathcal{R}[\phi_1] - id| \leq C|\phi_1|$. Estimating the third line requires no intermediate steps. \square

4 The exit manifold

4.1 Isolating the center directions

So long as U and V are not too far from the set of translations and rotations of Q in $W^{8/5,5}$ we can use Lemma 17 to select the collective coordinates $\mathbf{r}_j(t)$ and $\theta_j(t)$ so that

$$\Pi^c W_1(\cdot, t) = 0 \quad \text{and} \quad \Pi^c W_2(\cdot, t) = 0. \quad (11)$$

Applying Π^s to (8) and (9) we have, because of (11),

$$\begin{aligned} \partial_t W_1 &= A W_1 + \Pi^s B_1(\mathbf{m}) W_2 + \Pi^s J_1 \\ \partial_t W_2 &= A W_2 + \Pi^s B_2(\mathbf{m}) W_1 + \Pi^s J_2. \end{aligned} \quad (12)$$

We can isolate the equations of motion for \mathbf{r}_j and θ_j by applying Π^c to (8) and (9). If we coordinatize E^c as

$$\Pi^c (\mathbf{a} \cdot \nabla Q(\mathbf{y}) + \phi \mathbf{y}^\perp \cdot \nabla Q(\mathbf{y})) = \begin{pmatrix} \mathbf{a} \\ \phi \end{pmatrix} \in \mathbf{R}^3,$$

applying the projections gives:

$$\begin{aligned} \begin{pmatrix} \mathcal{R}[-\theta_1] \dot{\mathbf{r}}_1 - \mathbf{c}_1 \\ \dot{\theta}_1 \end{pmatrix} &= -\Pi^c (B_1(\mathbf{m}) W_2(\mathbf{y}_2, t) + K_1 + J_{1,mod} + J_{1,int} + J_{1,nl}) \\ \begin{pmatrix} \mathcal{R}[-\theta_2] \dot{\mathbf{r}}_2 - \mathbf{c}_2 \\ \dot{\theta}_2 \end{pmatrix} &= -\Pi^c (B_2(\mathbf{m}) W_1(\mathbf{y}_1, t) + K_2 + J_{2,mod} + J_{2,int} + J_{2,nl}). \end{aligned} \quad (13)$$

Since K_1 and K_2 depend explicitly on $\dot{\mathbf{r}}_j(t)$ and $\dot{\theta}_j(t)$, (13) does not quite constitute the equations of motion. The following lemma helps us isolate $\dot{\mathbf{r}}_j(t)$ and $\dot{\theta}_j(t)$:

Lemma 24.

$$\Pi^c K_j = L_j(\mathbf{m}, \nabla W_j) \begin{pmatrix} \mathcal{R}[-\theta_j] \dot{\mathbf{r}}_j - \mathbf{c}_j \\ \dot{\theta}_j \end{pmatrix} \quad (14)$$

and if $|\phi_j(t)| \leq \pi/2$ then:

$$\|L_j(\mathbf{m}(t), \nabla W_j(t))\| \leq C |\phi_j(t)| + C \|W_j(t)\|_{L^5}.$$

Proof. Notice that we can rewrite $\Pi^c K_1$ as:

$$\begin{aligned} \Pi^c K_1 &= \Pi^c ((\nabla Q(\mathbf{z}_1) - \nabla Q(\mathbf{y})) + \mathcal{R}[-\phi_1] \nabla W_1) \cdot \{\mathcal{R}[-\theta_1] \dot{\mathbf{r}}_1 - \mathbf{c}_1\} \\ &\quad + \Pi^c (\mathcal{R}[-\phi_1] \mathbf{y}^\perp \cdot (\nabla Q(\mathbf{z}_1) - \nabla Q(\mathbf{y})) + (\mathcal{R}[-\phi_1] \mathbf{y}^\perp - \mathbf{y}^\perp) \cdot \nabla Q(\mathbf{y})) \dot{\theta}_1 \end{aligned}$$

Thus we have (14). Now we estimate the various terms in $\Pi^c K_1$. We use the estimates in Lemma 23 to observe that the second line above can be bounded above by:

$$C |\phi_1(t)| |\dot{\theta}_1|.$$

Similarly, we saw in the proof of Lemma 23 that $\|\nabla Q(\mathbf{z}_1) - \nabla Q(\mathbf{y})\|_{L^2} \leq C |\phi_1(t)|$ and since we know from Hypothesis 2 that Π^c is computed by taking an L^2 inner product with the adjoint function, we have:

$$|\Pi^c (\nabla Q(\mathbf{z}_1) - \nabla Q(\mathbf{y})) \cdot \{\mathcal{R}[-\theta_1] \dot{\mathbf{r}}_1 - \mathbf{c}_1\}| \leq C |\phi_1(t)| |\mathcal{R}[-\theta_1] \dot{\mathbf{r}}_1 - \mathbf{c}_1|.$$

Finally, we have by an integration by parts and Hölder's inequality:

$$\begin{aligned} |\Pi^c(\mathcal{R}[-\phi_1]\nabla W_1) \cdot \{\mathcal{R}[-\theta_1]\dot{\mathbf{r}}_1 - \mathbf{c}\mathbf{i}\}| &= |\langle \psi^\dagger, [(\mathcal{R}[-\phi_1]\nabla W_1) \cdot \{\mathcal{R}[-\theta_1]\dot{\mathbf{r}}_1 - \mathbf{c}\mathbf{i}\}] \rangle| \\ &= |\langle [\mathcal{R}[\phi_1] (\mathcal{R}[-\theta_1]\dot{\mathbf{r}}_1 - \mathbf{c}\mathbf{i})] \cdot \nabla \psi^\dagger, W_1 \rangle| \\ &\leq |\mathcal{R}[-\theta_1]\dot{\mathbf{r}}_1 - \mathbf{c}\mathbf{i}| \|\nabla \psi^\dagger\|_{L^{5/4}} \|W_1\|_{L^5} \end{aligned}$$

Recall that Hypothesis 2 guarantees that $\|\nabla \psi^\dagger\|_{L^{5/4}} < \infty$. This completes the proof. \square

With this Lemma, provided \mathbf{m} and W are small, we have no problem inverting $id + L_j$. Therefore:

$$\begin{aligned} \begin{pmatrix} \mathcal{R}[-\theta_1]\dot{\mathbf{r}}_1 - \mathbf{c}\mathbf{i} \\ \dot{\theta}_1 \end{pmatrix} &= -\Pi^c (B_1(\mathbf{m})W_2(\mathbf{y}_2, t) + J_1 - K_1 + \check{K}_1) \\ \begin{pmatrix} \mathcal{R}[-\theta_2]\dot{\mathbf{r}}_2 - \mathbf{c}\mathbf{i} \\ \dot{\theta}_2 \end{pmatrix} &= -\Pi^c (B_2(\mathbf{m})W_1(\mathbf{y}_1, t) + J_2 - K_2 + \check{K}_2). \end{aligned}$$

where

$$\check{K}_1 = -((id + L_j)^{-1} - id) \Pi^c (B_1 W_2(\mathbf{y}_2, t) + J_1 - K_1)$$

and \check{K}_2 is defined analogously. Our previous estimates show that there is a constant $C > 0$ so that if $|\mathbf{m}|_{C^1[0, T_1]} \leq \pi/2$ then

$$|\check{K}_1(t)| \leq C (|\mathbf{m}(t)| + \|W_1\|_{L^5}) (\|W_2\|_{L^5} + \|J_1 - K_1\|_{L^5}). \quad (15)$$

Recalling their definitions, we see that the evolution equations for \mathbf{p}_j and ϕ_j are:

$$\begin{aligned} \begin{pmatrix} \dot{\mathbf{p}}_1 \\ \dot{\phi}_1 \end{pmatrix} &= -\tilde{\mathcal{R}}[\phi_1 + \theta_{10}] \Pi^c (B_1 W_2(\mathbf{y}_2, t) + J_1 - K_1 + \check{K}_1) \\ \begin{pmatrix} \dot{\mathbf{p}}_2 \\ \dot{\phi}_2 \end{pmatrix} &= -\tilde{\mathcal{R}}[\phi_2 + \theta_{20}] \Pi^c (B_2 W_1(\mathbf{y}_1, t) + J_2 - K_2 + \check{K}_2) \end{aligned} \quad (16)$$

with

$$\tilde{\mathcal{R}}[\theta] := \begin{bmatrix} \mathcal{R}[\theta] & 0 \\ 0 & 1 \end{bmatrix}.$$

Equations (12) and (16) are equivalent to (2), and our goal now is to solve this system. To compress notation, we let

$$\begin{aligned} \mathbf{W} &= (W_1, W_2) \\ \mathbf{m} &= (\mathbf{p}_1, \phi_1, \mathbf{p}_2, \phi_2) \\ \mathbf{J}_s(\mathbf{m}, \mathbf{W}) &= (J_1, J_2) \\ \mathbf{J}_{cc}(\mathbf{m}, \mathbf{W}) &= (J_1 - K_1, J_2 - K_2) \\ \mathbf{K}_{cc}(\mathbf{m}, \mathbf{W}) &= (\check{K}_1, \check{K}_2) \\ \mathbf{A} &= \begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix} \\ \mathbf{B}(\mathbf{m}) &= \begin{bmatrix} 0 & B_1 \\ B_2 & 0 \end{bmatrix} \\ \mathbf{P}^s &:= \text{diag}(\Pi^s, \Pi^s), \quad \mathbf{P}^c := \text{diag}(\Pi^c, \Pi^c) \end{aligned}$$

$$\mathbf{R}(\mathbf{m}) := \begin{bmatrix} \tilde{\mathcal{R}}[\phi_1 + \theta_{10}] & 0 \\ 0 & \tilde{\mathcal{R}}[\phi_2 + \theta_{20}] \end{bmatrix}$$

and then we see that (12) and (16) become:

$$\begin{aligned} \mathbf{W}_t &= \mathbf{A}\mathbf{W} + \mathbf{P}^s(\mathbf{B}(\mathbf{m})\mathbf{W} + \mathbf{J}_s(\mathbf{m}, \mathbf{W})) \\ \dot{\mathbf{m}} &= -\mathbf{R}(\mathbf{m})\mathbf{P}^c(\mathbf{B}(\mathbf{m})\mathbf{W} + \mathbf{J}_{cc}(\mathbf{m}, \mathbf{W})) - \mathbf{R}(\mathbf{m})\mathbf{K}_{cc}(\mathbf{m}, \mathbf{W}) \end{aligned} \quad (17)$$

Note that if we take the appropriate $L^5[0, T]$ norms of the estimates in Lemma 20-23 we arrive at the following estimates for \mathbf{J}_s and \mathbf{J}_{cc} :

Lemma 25. *There is a constant $C > 0$ so that if $\|\mathbf{m}\|_{C^1[0, T]} \leq \pi/2$:*

$$\begin{aligned} \|\mathbf{J}_s\|_{X_{a,0}[T]} &\leq C \left\{ \|\mathbf{m}\|_{C^1([0, T])}^2 + \|\mathbf{m}\|_{C^1([0, T])} \|\mathbf{W}\|_{X'_{a,0}[T]} \right. \\ &\quad \left. + N(\mathbf{m}) + e^{-\frac{b_0}{2}\mu_0} \|\mathbf{W}\|_{X'_{a,0}[T]} + \|\mathbf{W}\|_{X'_{a,0}[T]}^2 \right\}, \end{aligned} \quad (18)$$

and

$$\|\mathbf{J}_{cc}\|_{X_{a,0}[T]} \leq C \left\{ \|\mathbf{m}\|_{C^1([0, T])} \|\mathbf{W}\|_{X'_{a,0}[T]} + N(\mathbf{m}) + e^{-\frac{b_0}{2}\mu_0} \|\mathbf{W}\|_{X'_{a,0}[T]} + \|\mathbf{W}\|_{X'_{a,0}[T]}^2 \right\}. \quad (19)$$

Here

$$N(\mathbf{m}) := \left(\int_0^T \left| e^{at} e^{-\frac{b_0}{2}\mu(t)} \right|^5 dt \right)^{1/5} = \|\exp(a \diamond -b_0/2\mu(\diamond))\|_{L^5[0, T]}.$$

The constant C does not depend on $0 \leq T \leq \infty$.

4.2 Stability of exits

We now can Theorem 5. We must first show that (given appropriate initial pulse positions and orientations) smallness of the modulation parameters $\mathbf{m}(t)$ implies that $\mu(t)$ is large. This in turn implies that \mathbf{J}_s , \mathbf{J}_{cc} and \mathbf{K}_{cc} will be small. Recall that $\mu^*(t)$ is the separation distance between pulses if $\mathbf{m} = 0$ and that μ_0^* is the minimum separation distance in this situation. We need:

Lemma 26. *For all $\Delta\mathbf{v}_0 \neq 0$ and $\Delta\mathbf{r}_0$ with $\mu_0^* > 0$, there exists $\delta_1 > 0$ so that $|\mathbf{m}(t)| \leq \delta_1$ (for $t \geq 0$) implies*

$$\mu(t) \geq \frac{1}{2}\mu^*(t).$$

for all $t \geq 0$. Additionally, with $k^* := c|\Delta\mathbf{v}_0| \frac{\mu_0^*}{|\Delta\mathbf{r}_0|}$,

$$N(\mathbf{m}) = \left| \int_0^\infty (e^{\frac{b_0}{8}k^*t} e^{-\frac{b_0}{2}\mu(t)})^5 dt \right|^{1/5} \leq C|\Delta\mathbf{v}_0|^{-1/5} e^{-\frac{b_0}{4}\mu_0^*}.$$

Proof. (Lemma 26) Recall from the Introduction that

$$\mu^*(t) \geq k^*t$$

where k^* is as in the statement of the lemma.

Take

$$\delta_1 = \min \left\{ \frac{\mu_0^*}{4}, \frac{k^*}{8c} \right\}.$$

The definition of \mathbf{p}_j and ϕ_j together with the triangle inequality give

$$\begin{aligned}
\mu(t) &\geq \mu^*(t) - \sum_{j=1}^2 \left(|\mathbf{p}_j(t)| + c \left| \int_0^t [\mathcal{R}[\phi_j(s) + \theta_{j0}] - \mathcal{R}[\theta_{j0}]] \mathbf{i} ds \right| \right) \\
&\geq \mu^*(t) - \sum_{j=1}^2 \left(|\mathbf{p}_j(t)| + ct \sup_{0 \leq s \leq t} |\mathcal{R}[\phi_j(s) + \theta_{j0}] - \mathcal{R}[\theta_{j0}]]| \right) \\
&\geq \mu^*(t) - \delta_1 - 2ct\delta_1 \\
&\geq \frac{1}{2}\mu^*(t) + \frac{1}{4}\mu_0^* - \delta_1 + \frac{1}{4}(\mu^*(t) - 8ct\delta_1) \\
&\geq \frac{1}{2}\mu^*(t).
\end{aligned}$$

The integral estimate follows from this. Here is the calculation:

$$\begin{aligned}
\int_0^\infty (e^{\frac{b_0}{8}k^*t} e^{-\frac{b_0}{2}\mu(t)})^5 dt &\leq e^{-\frac{5b_0}{4}\mu_0^*} \int_0^\infty e^{\frac{5b_0}{8}k^*t - \frac{5b_0}{4}\mu^*(t)} dt \\
&\leq e^{-\frac{5b_0}{4}\mu_0^*} \int_0^\infty e^{-\frac{5b_0}{8}k^*t} dt \\
&\leq C|\Delta\mathbf{v}_0|^{-1} e^{-\frac{5b_0}{4}\mu_0^*}.
\end{aligned}$$

□

Proof. (Theorem 5) If the operator \mathbf{B} was small in norm on $X_{a,0}$, we could use this and a straightforward perturbation argument to prove our result. Instead we must make one change to (17), which is motivated by (4). Consider the system:

$$\begin{aligned}
\mathbf{W}_t^{ic} &= \mathbf{A}\mathbf{W}^{ic} + \mathbf{P}^s \mathbf{J}_s(\mathbf{m}, \mathbf{W}^{ic} + \mathbf{W}^{loc}) \\
\mathbf{W}_t^{loc} &= \mathbf{A}\mathbf{W}^{loc} + \mathbf{P}^s (\mathbf{B}(\mathbf{m}) (\mathbf{W}^{ic} + \mathbf{W}^{loc})) \\
\dot{\mathbf{m}} &= -\mathbf{R}(\mathbf{m})\mathbf{P}^c (\mathbf{B}(\mathbf{m}) (\mathbf{W}^{ic} + \mathbf{W}^{loc})) + \mathbf{J}_{cc}(\mathbf{m}, \mathbf{W}^{ic} + \mathbf{W}^{loc}) \\
&\quad - \mathbf{R}(\mathbf{m})\mathbf{K}_{cc}(\mathbf{m}, \mathbf{W}^{ic} + \mathbf{W}^{loc})
\end{aligned} \tag{20}$$

with initial conditions

$$\mathbf{W}^{ic}(\mathbf{y}, 0) = \mathbf{W}_0 \in (E^s)^2 \cap W^{8/5,5}, \quad \mathbf{W}^{loc}(\mathbf{y}, 0) = 0 \quad \text{and} \quad \mathbf{m}(0) = 0.$$

If $\mathbf{W} = \mathbf{W}^{ic} + \mathbf{W}^{loc}$ then \mathbf{W} solves (17). Notice that right hand side of the equation for \mathbf{W}^{ic} consists of terms which are small in $X_{a,0}[T]$ due to Lemma 25 and estimate (15). The right hand side of the equation for \mathbf{W}^{loc} is in $X_{a,b_0}[T]$ even if $\mathbf{W}^{ic} \in X_{a,0}[T]$, due to the localizing property of \mathbf{B} seen in Lemma 19. (The superscript “ic” represents “initial condition” and the superscript “loc” stands for “localized”.)

In Theorem 27 in the Appendix we establish that $\|\mathbf{W}_0\|_{W^{8/5,5}} \leq \delta_2$ implies that (20) has a unique solution

$$(\mathbf{W}^{ic}, \mathbf{W}^{loc}, \mathbf{m}) \in X'_{a,0}[T_1] \times X'_{a,b_0}[T_1] \times C^1[0, T_1]$$

for some $0 < T_1 \leq \infty$, where we have taken $a = \min\{\alpha/2, b_0k^*/8\}$. $\delta_2 > 0$ is a universal

constant. Such a solution satisfies the integral equation:

$$\begin{aligned}
\mathbf{W}^{ic}(t) &= e^{\mathbf{A}t} \mathbf{W}_0 + \int_0^t e^{\mathbf{A}(t-\tau)} \mathbf{P}^s \mathbf{J}_s(\mathbf{m}(\tau), \mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau)) d\tau \\
\mathbf{W}^{loc}(t) &= \int_0^t e^{\mathbf{A}(t-\tau)} \mathbf{P}^s \mathbf{B}(\mathbf{m}(\tau)) (\mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau)) d\tau \\
\mathbf{m}(t) &= - \int_0^t \left\{ \mathbf{R}(\mathbf{m}(\tau)) \mathbf{P}^c (\mathbf{B}(\mathbf{m}(\tau)) (\mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau)) \right. \\
&\quad \left. + \mathbf{J}_{cc}(\mathbf{m}(\tau), \mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau))) + \mathbf{R}[\mathbf{m}(\tau)] \mathbf{K}_{cc}(\mathbf{m}(\tau), \mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau)) \right\} d\tau.
\end{aligned} \tag{21}$$

Let

$$\mathcal{K}(T) := \|\mathbf{W}^{ic}\|_{X'_{a,0}[T]} + \|\mathbf{W}^{loc}\|_{X'_{a,b_0}[T]} + \|\mathbf{m}\|_{C^1[0,T]}.$$

Here, $0 < T \leq T_1$ and we assume that $\mathcal{K}(T) \leq \delta_1$, which allows us to apply Lemma 26 when needed.

Applying Theorem 15 to (21) tells us that

$$\begin{aligned}
\|\mathbf{W}^{ic}\|_{X'_{a,0}[T]} &\leq C \left(\|\mathbf{W}_0\|_{W^{8/5,5}} + \|\mathbf{J}_s(\mathbf{m}, \mathbf{W}^{ic} + \mathbf{W}^{loc})\|_{X_{a,0}[T]} \right) \\
\|\mathbf{W}^{loc}\|_{X'_{a,b_0}[T]} &\leq C \|\mathbf{B}(\mathbf{m}) (\mathbf{W}^{ic} + \mathbf{W}^{loc})\|_{X_{a,b_0}[T]}.
\end{aligned} \tag{22}$$

The estimates in Lemma 25 and Lemma 26 yield:

$$\begin{aligned}
\|\mathbf{W}^{ic}\|_{X'_{a,0}[T]} &\leq C \left\{ \|\mathbf{W}_0\|_{W^{8/5,5}} + \|\mathbf{m}\|_{C^1([0,T])}^2 \right. \\
&\quad \left. + \left(\|\mathbf{m}\|_{C^1([0,T])} + e^{-\frac{b_0}{4}\mu_0^*} \right) \left(\|\mathbf{W}^{ic}\|_{X'_{a,0}[T]} + \|\mathbf{W}^{loc}\|_{X'_{a,b_0}[T]} \right) \right. \\
&\quad \left. + C |\Delta \mathbf{v}_0|^{-1/5} e^{-\frac{b_0}{4}\mu_0^*} + \|\mathbf{W}^{ic}\|_{X'_{a,0}[T]}^2 + \|\mathbf{W}^{loc}\|_{X'_{a,b_0}[T]}^2 \right\} \\
&\leq C_1^* |\Delta \mathbf{v}_0|^{-1/5} \left(\|\mathbf{W}_0\|_{W^{8/5,5}} + e^{-\frac{b_0}{4}\mu_0^*} + e^{-\frac{b_0}{4}\mu_0^*} \mathcal{K}(T) + \mathcal{K}^2(T) \right).
\end{aligned} \tag{23}$$

Note that the constant $C_1^* > 0$ does not depend upon $\mathcal{K}(T)$, T , a , b_0 or $\|\mathbf{W}_0\|_{W^{8/5,5}}$.

Similarly if we apply Lemma 19 we get

$$\|\mathbf{W}^{loc}\|_{X'_{a,b_0}[T]} \leq C \left\{ e^{-\frac{b_0}{4}\mu_0^*} \|\mathbf{W}^{loc}\|_{X'_{a,b_0}[T]} + \|\mathbf{W}^{ic}\|_{X'_{a,0}[T_1]} \right\} \tag{24}$$

which if we combine with the last inequality for $\|\mathbf{W}^{ic}\|_{X'_{a,0}[T]}$ we get

$$\|\mathbf{W}^{loc}\|_{X'_{a,b_0}[T]} \leq C_2^* |\Delta \mathbf{v}_0|^{-1/5} \left\{ \|\mathbf{W}_0\|_{W^{8/5,5}} + e^{-\frac{b_0}{4}\mu_0^*} + e^{-\frac{b_0}{4}\mu_0^*} \mathcal{K}(T) + \mathcal{K}^2(T) \right\}. \tag{25}$$

$C_2^* > 0$ does not depend upon $\mathcal{K}(T)$, T , a , b_0 or $\|\mathbf{W}_0\|_{W^{8/5,5}}$.

We wish to estimate \mathbf{m} in $C^1[0, T]$. We begin with:

$$\begin{aligned}
\|\mathbf{m}\|_{C[0,T]} &\leq C \sup_{0 \leq t \leq T} \int_0^t \left\{ \|\mathbf{B}(\mathbf{m}(\tau)) (\mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau))\|_{L^5} \right. \\
&\quad \left. + \|\mathbf{J}_{cc}(\mathbf{m}(\tau), \mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau))\|_{L^5} + |\mathbf{K}_{cc}(\mathbf{m}(\tau), \mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau))| \right\} d\tau.
\end{aligned} \tag{26}$$

To estimate (26) we first apply Lemma 19:

$$|\mathbf{m}(t)| \leq C \sup_{0 \leq t \leq T} \int_0^t \left\{ \|\mathbf{W}^{ic}(\tau)\|_{L^5} + e^{-\frac{b_0}{4}\mu_0^*} \|\mathbf{W}^{loc}(\tau)\|_{L_{b_0}^5} \right. \\ \left. + \|\mathbf{J}_{cc}(\mathbf{m}(\tau), \mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau))\|_{L^5} + |\mathbf{K}_{cc}(\mathbf{m}(\tau), \mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau))| \right\} d\tau.$$

We want a bound independent of t . We have by Hölder's inequality for a general function \mathbf{F} and $0 \leq b \leq b_0$:

$$\int_0^t \|\mathbf{F}(\tau)\|_{L_b^5} d\tau = \int_0^t e^{-b_0 k^* \tau/8} e^{b_0 k^* \tau/8} \|\mathbf{F}(\tau)\|_{L_b^5} d\tau \\ \leq |b_0 k^*/8|^{-4/5} \|\mathbf{F}\|_{X_{a,b}[T]} \\ \leq C |\Delta \mathbf{v}_0|^{-4/5} \|\mathbf{F}\|_{X_{a,b}[T]}.$$
(27)

Applying this to all but the \mathbf{K}_{cc} term and we have

$$|\mathbf{m}(t)| \leq C |\Delta \mathbf{v}_0|^{-4/5} \left\{ \|\mathbf{W}^{ic}\|_{X_{a,0}[T]} + e^{-\frac{b_0}{4}\mu_0^*} \|\mathbf{W}^{loc}\|_{X_{a,b_0}[T]} + \|\mathbf{J}_{cc}(\mathbf{m}, \mathbf{W}^{ic} + \mathbf{W}^{loc})\|_{X_{a,0}[T]} \right\} \\ + \sup_{0 \leq t \leq T} \int_0^t |\mathbf{K}_{cc}(\mathbf{m}(\tau), \mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau))| d\tau$$

Using (19), Lemma 26 and the estimate (23) tell us that the first line above is bounded by:

$$C |\Delta \mathbf{v}_0|^{-1} \left\{ \|\mathbf{W}_0\|_{W^{8/5,5}} + e^{-\frac{b_0}{4}\mu_0^*} + e^{-\frac{b_0}{4}\mu_0^*} \mathcal{K}(T) + \mathcal{K}^2(T) \right\}.$$

For the second line, applying (15) gives

$$INT := \sup_{0 \leq t \leq T} \int_0^t |\mathbf{K}_{cc}(\mathbf{m}(\tau), \mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau))| d\tau \\ \leq C \sup_{0 \leq t \leq T} \int_0^t \left(\|\mathbf{m}\|_{C[0,T]} + \|\mathbf{W}^{ic}\|_{L^5} + \|\mathbf{W}^{loc}\|_{L_{b_0}^5} \right) \\ \left(\|\mathbf{W}^{ic}\|_{L^5} + \|\mathbf{W}^{loc}\|_{L_{b_0}^5} + \|\mathbf{J}_{cc}\|_{L_{b_0}^5} \right) d\tau.$$

The embedding (6) and (27) then imply

$$INT \leq C |\Delta \mathbf{v}_0|^{-4/5} \left(\|\mathbf{m}\|_{C[0,T]} + \|\mathbf{W}^{ic}\|_{X'_{a,0}[T]} + \|\mathbf{W}^{loc}\|_{X'_{a,b_0}[T]} \right) \\ \left(\|\mathbf{W}^{ic}\|_{X_{a,0}[T]} + \|\mathbf{W}^{loc}\|_{X_{a,b_0}[T]} + \|\mathbf{J}_{cc}\|_{X_{a,0}[T]} \right)$$

We consolidate the estimate for the two pieces to get:

$$\|\mathbf{m}\|_{C[0,T]} \leq C_3^* |\Delta \mathbf{v}_0|^{-1} (1 + \mathcal{K}(T)) \left\{ \|\mathbf{W}_0\|_{W^{8/5,5}} + e^{-\frac{b_0}{4}\mu_0^*} + e^{-\frac{b_0}{4}\mu_0^*} \mathcal{K}(T) + \mathcal{K}^2(T) \right\}.$$

The constant $C_3^* > 0$ does not depend upon $\mathcal{K}(T)$, T , a , b_0 or $\|\mathbf{W}_0\|_{W^{8/5,5}}$.

We must also estimate $\|\dot{\mathbf{m}}\|_{C[0,T]}$. Since $\|\mathbf{P}^s \mathbf{F}\| \leq C \|\mathbf{F}\|_{L^5}$ it is clear from (20) and (6) that

$$\|\dot{\mathbf{m}}\|_{C[0,T]} \leq C \sup_{t \in [0,T]} \left\{ \|\mathbf{W}^{ic}\|_{L^5} + e^{-\frac{b_0}{4}\mu_0^*} \|\mathbf{W}^{loc}\|_{L^5} + \|\mathbf{J}_{cc}\|_{L^5} + |\mathbf{K}_{cc}(t)| \right\} \\ \leq C \left\{ \|\mathbf{W}^{ic}\|_{X'_{a,0}[T]} + e^{-\frac{b_0}{4}\mu_0^*} \|\mathbf{W}^{loc}\|_{X'_{a,b_0}[T]} + \|\mathbf{J}_{cc}\|_{X'_{a,0}[T]} + \sup_{t \in [0,T]} |\mathbf{K}_{cc}(t)| \right\} \quad (28)$$

With the exception of $\sup_{t \in [0, T]} |\mathbf{K}_{cc}(t)|$ we have estimated each of these pieces already. This last piece can be estimated much like INT and we omit the details as they are in fact simpler than those for INT .

Therefore

$$\|\mathbf{m}\|_{C^1[0, T]} \leq C_4^* |\Delta \mathbf{v}_0|^{-1} (1 + \mathcal{K}(T)) \left\{ \|\mathbf{W}_0\|_{W^{8/5, 5}} + e^{-\frac{b_0}{4} \mu_0^*} + e^{-\frac{b_0}{4} \mu_0^*} \mathcal{K}(T) + \mathcal{K}^2(T) \right\}. \quad (29)$$

The constant $C_4^* > 0$ does not depend upon $\mathcal{K}(T)$, T , a , b_0 or $\|\mathbf{W}_0\|_{W^{8/5, 5}}$.

Putting (23), (25) and (29) together gives:

$$\mathcal{K}(T) \leq C^* |\Delta \mathbf{v}_0|^{-1} (1 + \mathcal{K}(T)) \left\{ \|\mathbf{W}_0\|_{W^{8/5, 5}} + e^{-\frac{b_0}{4} \mu_0^*} + e^{-\frac{b_0}{4} \mu_0^*} \mathcal{K}(T) + \mathcal{K}^2(T) \right\}. \quad (30)$$

where $C^* = \max \{C_1^*, C_2^*, C_4^*\}$

Let

$$\mathcal{K}_0 := \min \left\{ 1, \delta_1, \tilde{C} \delta_2, \frac{|\Delta \mathbf{v}_0|}{8C^*} \right\},$$

$$\delta_0 := \frac{\mathcal{K}_0 |\Delta \mathbf{v}_0|}{4C^* (1 + \mathcal{K}_0)}$$

and

$$\epsilon_{20} = \frac{\mathcal{K}_0 |\Delta \mathbf{v}_0|}{4C^* (1 + \mathcal{K}_0)^2}$$

(Here, δ_1 is as in Lemma 26, δ_2 is as in Theorem 27 and $1/\tilde{C}$ is the constant from the embedding (6).) Suppose that

$$\|\mathbf{W}_0\|_{W^{8/5, 5}} \leq \delta_{exit}$$

and

$$\mu_0^* \geq -\frac{4}{b_0} \ln(\epsilon_{20}) =: M_0.$$

Also set

$$T_2 := \sup \{T \leq T_1 : \mathcal{K}(t) \leq \mathcal{K}_0 \text{ for all } 0 \leq t \leq T\}.$$

Since $\mathcal{K}(0) = 0$ this is either nonnegative or infinite. If it is infinite then we have proven our main result. If it is finite then we must have $\mathcal{K}(T_2) = \mathcal{K}_0$. Otherwise, Theorem 27 and (6) imply that the solution exists for some slightly longer time and moreover $\mathcal{K}(t)$ is continuous on this larger interval.

However, (30) and our choices for \mathcal{K}_0 , δ_0 and ϵ_{20} imply:

$$C^* |\Delta \mathbf{v}_0|^{-1} (1 + \mathcal{K}) \mathcal{K}^2 \leq 2C^* |\Delta \mathbf{v}_0|^{-1} \mathcal{K}_0^2 \leq \frac{\mathcal{K}_0}{4}$$

$$C^* |\Delta \mathbf{v}_0|^{-1} (1 + \mathcal{K}) \|\mathbf{W}_0\|_{W^{8/5, 5}} \leq C^* |\Delta \mathbf{v}_0|^{-1} (1 + \mathcal{K}_0) \delta_0 \leq \frac{\mathcal{K}_0}{4}$$

and

$$C^* |\Delta \mathbf{v}_0|^{-1} (1 + \mathcal{K})^2 e^{-\frac{b_0}{4} \mu_0^*} \leq C^* |\Delta \mathbf{v}_0|^{-1} (1 + \mathcal{K}_0)^2 \epsilon_{20} \leq \frac{\mathcal{K}_0}{4}.$$

which all together imply

$$\mathcal{K}_0 \leq 3/4 \mathcal{K}_0$$

which is impossible. Therefore T_2 is not finite and we are done. \square

5 The shooting manifold

5.1 Existence of shooting solutions

We now prove the existence of the shooting solution described in Theorem 7

Proof. (Theorem 7) Recall from Section 3 that for an arbitrary choice \mathbf{m} , equations (8)-(9) describe the evolution of $\mathbf{W} = (W_1, W_2)$. In the previous sections it was desirable to choose \mathbf{m} so that $\mathbf{W} \in E^s \times E^s$ for all $t > 0$. Now it is useful to set $\mathbf{m} = 0$, which is to say we take $\mathbf{r}_j(t) = \mathbf{r}_j^*(t) = \mathbf{r}_{j0} + ct\mathbf{v}(\theta_{j0})$ and $\theta_j(t) = \theta_{j0}$. We make the assumption that $\Delta\mathbf{r}_0 \cdot \Delta\mathbf{v}_0 \leq 0$ which implies that

$$\mu(t) \geq \max\{|\Delta\mathbf{r}_0|, -c|\Delta\mathbf{v}_0|t\}$$

for $t \leq 0$.

In this case (2) becomes:

$$\begin{aligned}\partial_t W_1 &= AW_1 + B_1(0)W_2 + J_{1,int} + J_{1,nl} \\ \partial_t W_2 &= AW_2 + B_2(0)W_1 + J_{2,int} + J_{2,nl}\end{aligned}$$

which we write compactly as

$$\mathbf{W}_t = \mathbf{A}\mathbf{W} + \mathbf{B}(0)\mathbf{W} + \mathbf{J}_{shoot}(\mathbf{W}). \quad (31)$$

Let $\eta = -b_0c|\Delta\mathbf{v}_0|/8$ and consider the map $\Psi : Z'_{\eta, b_0} \rightarrow Z'_{\eta, b_0}$ defined by

$$\Psi(\mathbf{W})(t) = \int_{-\infty}^t e^{\mathbf{A}(t-\tau)} (\mathbf{B}(0)\mathbf{W}(\tau) + \mathbf{J}_{shoot}(\mathbf{W}(\tau))) d\tau.$$

We prove that there exists $\mathcal{K}_1 > 0$ with property that $\|\mathbf{W}\|_{Z_{\eta, b_0}} \leq \mathcal{K}_1$ implies

$$\|\Psi(\mathbf{W})\|_{Z'_{\eta, b_0}} \leq \mathcal{K}_1 \quad (32)$$

and

$$\left\| \Psi(\mathbf{W}) - \Psi(\tilde{\mathbf{W}}) \right\|_{Z'_{\eta, b_0}} \leq \frac{1}{2} \left\| \mathbf{W} - \tilde{\mathbf{W}} \right\|_{Z'_{\eta, b_0}} \quad (33)$$

This implies Ψ has a unique fixed point $\mathbf{W}^* = (W_1^*, W_2^*) \in Z'_{\eta, b_0}$ which solves (31). These are the functions described in Theorem 7, and once we establish their existence, the proof is complete.

To prove the estimate (32) we apply Theorem 15

$$\|\Psi(\mathbf{W})\|_{Z'_{\eta, b_0}} \leq C \left(\|\mathbf{B}(0)\mathbf{W}\|_{Z_{\eta, b_0}} + \|\mathbf{J}_{shoot}(\mathbf{W})\|_{Z_{\eta, b_0}} \right).$$

Lemma 19 implies

$$\|\mathbf{B}(0)\mathbf{W}\|_{Z_{\eta, b_0}} \leq C e^{-\frac{b_0}{2}|\Delta\mathbf{r}_0|} \|\mathbf{W}\|_{Z_{\eta, b_0}}.$$

Lemmas 21 and 22 imply

$$\|\mathbf{J}_{shoot}\|_{Z_{\eta, b_0}} \leq C \left(N + e^{-\frac{b_0}{2}|\Delta\mathbf{r}_0|} \|\mathbf{W}\|_{Z_{\eta, b_0}} + \|\mathbf{W}\|_{Z_{\eta, b_0}}^2 \right)$$

where

$$\begin{aligned}N &= \left(\int_{\mathbf{R}^-} \left| e^{-\frac{b_0}{2}\mu(t)} e^{\eta t} \right|^5 dt \right)^{1/5} \leq e^{-\frac{b_0}{4}|\Delta\mathbf{r}_0|} \left(\int_{\mathbf{R}^-} \left| e^{\frac{b_0}{4}c|\Delta\mathbf{v}_0|t} e^{\eta t} \right|^5 dt \right)^{1/5} \\ &\leq e^{-\frac{b_0}{4}|\Delta\mathbf{r}_0|} \left(\int_{\mathbf{R}^-} \left| e^{\frac{b_0}{8}c|\Delta\mathbf{v}_0|t} \right|^5 dt \right)^{1/5} \leq C |\Delta\mathbf{v}_0|^{-1/5} e^{-\frac{b_0}{4}|\Delta\mathbf{r}_0|}.\end{aligned}$$

Therefore

$$\|\Psi(\mathbf{W})\|_{Z'_{\eta,b_0}} \leq C^* |\Delta \mathbf{v}_0|^{-1/5} \left(e^{-\frac{b_0}{4} |\Delta \mathbf{r}_0|} + e^{-\frac{b_0}{4} |\Delta \mathbf{r}_0|} \|\mathbf{W}\|_{Z'_{\eta,b_0}} + \|\mathbf{W}\|_{Z'_{\eta,b_0}}^2 \right).$$

Similarly,

$$\begin{aligned} \|\Psi(\mathbf{W}) - \Psi(\tilde{\mathbf{W}})\|_{Z'_{\eta,b_0}} &\leq C \left(\|\mathbf{B}(0)(\mathbf{W} - \tilde{\mathbf{W}})\|_{Z_{\eta,b_0}} + \|\mathbf{J}_{shoot}(\mathbf{W}) - \mathbf{J}_{shoot}(\tilde{\mathbf{W}})\|_{Z_{\eta,b_0}} \right) \\ &\leq C \left(e^{-\frac{b_0}{2} |\Delta \mathbf{r}_0|} \|\mathbf{W} - \tilde{\mathbf{W}}\|_{Z_{\eta,b_0}} + \sum_j \|J_{j,nl}(\mathbf{W}) - J_{j,nl}(\tilde{\mathbf{W}})\|_{Z_{\eta,b_0}} \right). \end{aligned}$$

To estimate the $J_{j,nl}$ terms, recall that F is C^2 so we have the following pointwise estimate for $J = F(Q + W) - F(Q) - F'(Q)W$ and $\tilde{J} = F(Q + \tilde{W}) - F(Q) - F'(Q)\tilde{W}$. (Here $Q, W, \tilde{W} \in \mathbf{R}^N$.) First $F(Q + W) - F(Q + \tilde{W}) = F'(Q + W_\gamma)(W - \tilde{W})$ where W_γ lies on the line segment connecting W to \tilde{W} . Then

$$\begin{aligned} |J - \tilde{J}| &= |F'(Q + W_\gamma)(W - \tilde{W}) - F'(Q)(W - \tilde{W})| \\ &\leq C |W_\gamma| |W - \tilde{W}| \\ &\leq C (|W| + |\tilde{W}|) |W - \tilde{W}| \end{aligned}$$

This yields

$$\|J_{j,nl}(\mathbf{W}) - J_{j,nl}(\tilde{\mathbf{W}})\|_{Z_{\eta,b_0}} \leq C \left(\|\mathbf{W}\|_{Z'_{\eta,b_0}} + \|\tilde{\mathbf{W}}\|_{Z'_{\eta,b_0}} \right) \|\mathbf{W} - \tilde{\mathbf{W}}\|_{Z'_{\eta,b_0}}.$$

Therefore

$$\|\Psi(\mathbf{W}) - \Psi(\tilde{\mathbf{W}})\|_{Z'_{\eta,b_0}} \leq C^{**} \left(e^{-\frac{b_0}{2} |\Delta \mathbf{r}_0|} + \|\mathbf{W}\|_{Z'_{\eta,b_0}} + \|\tilde{\mathbf{W}}\|_{Z'_{\eta,b_0}} \right) \|\mathbf{W} - \tilde{\mathbf{W}}\|_{Z_{\eta,b_0}} \quad (34)$$

Let

$$\mathcal{K}_1 = \min \left\{ \frac{|\Delta \mathbf{v}_0|^{1/5}}{3C^*}, \frac{1}{4C^{**}} \right\}$$

and

$$\epsilon_{10} = \min \left\{ \frac{\mathcal{K}_1 |\Delta \mathbf{v}_0|^{1/5}}{3C^*(1 + \mathcal{K}_1)}, \frac{1}{4C^{**}} \right\}.$$

If

$$\|\mathbf{W}\|_{Z'_{\eta,b_0}} \leq \mathcal{K}_1 \quad \text{and} \quad |\Delta \mathbf{r}_0| \geq -\frac{4}{b_0} \ln(\epsilon_{10}) =: M_1$$

then we have (32) and (33). We are done. \square

Before we move on to the stability of the shooting solution, we make a few remarks about the dependence of the functions \mathbf{W}^* on the trajectory parameters $\mathbf{n} = (\mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20})$. We claim that there exists a constant $C > 0$ so that

$$\|D_{\mathbf{n}} \mathbf{W}^*(-T)\|_{W^{8/5,5}} \leq C e^{\eta T/2}. \quad (35)$$

Notice that

$$D_{\mathbf{n}}\Psi(\mathbf{W})(t) = \int_{-\infty}^t e^{\mathbf{A}(t-\tau)} (D_{\mathbf{n}}\mathbf{B}(0)\mathbf{W}(\tau) + D_{\mathbf{n}}\mathbf{J}_{shoot}(\mathbf{W}(\tau))) d\tau$$

implies

$$\|D_{\mathbf{n}}\mathbf{W}^*\|_{Z'_{\eta/2,0}} \leq C \|D_{\mathbf{n}}\mathbf{B}(0)\mathbf{W}^*\|_{Z_{\eta/2,0}} + C \|D_{\mathbf{n}}\mathbf{J}_{shoot}(\mathbf{W}^*)\|_{Z_{\eta/2,0}}.$$

by Theorem 15 and (6). Provided we can show that the right hand side of this is bounded by a constant C , independent of \mathbf{n} , the embedding (6) will yield the claim. There is one minor complication, which is that $D_{\mathbf{n}}\mathbf{B}(0)\mathbf{W}^*$ contains terms like

$$\begin{aligned} & |\chi(\mathbf{x})D_{\theta_{10}} [(F'(Q(\mathbf{y}) + Q(\mathbf{y}_2)) - F'(Q(\mathbf{y}))W_2^*(\mathbf{y}_2))] \\ & \leq |F''(Q(\mathbf{y}) + Q(\mathbf{y}_2))| |\nabla Q(\mathbf{y}_2)| |D_{\theta_{10}}\mathbf{y}_2| |W_2^*(\mathbf{y}_2)| + |D_{\theta_{10}}\mathbf{y}_2| |\nabla W_2^*(\mathbf{y}_2)|. \end{aligned}$$

From Remark 18

$$|D_{\theta_{10}}\mathbf{y}_2| \leq (|D\mathcal{R}[\theta_{10} - \theta_{20}]\mathbf{y}| + |\mathbf{r}_2^*(t)| + ct |D\mathcal{R}[\theta_{10}]\mathbf{i}|) \leq C(1 + |\mathbf{y}_2| + t).$$

Likewise, we can control the derivatives with respect to \mathbf{r}_{10} , \mathbf{r}_{20} and θ_{20} . We conclude:

$$\|D_{\mathbf{n}}\mathbf{B}(0)\mathbf{W}^*(t)\|_{L^5} \leq C \|(1 + |\cdot| + t)\mathbf{W}^*(\cdot, t)\|_{W^{8/5,5}} \leq C \|(1 + t)\mathbf{W}^*(\cdot, t)\|_{W^{8/5,5}_{b_0}}$$

The secular growth here seems problematic, however $te^{-\eta t/2}$ is bounded for $t \leq 0$, therefore

$$\|D_{\mathbf{n}}\mathbf{B}(0)\mathbf{W}^*\|_{Z_{\eta/2,0}} \leq C \|\mathbf{W}^*\|_{Z'_{\eta,b_0}}.$$

In exactly the same way we can show

$$\|D_{\mathbf{n}}\mathbf{J}_{shoot}(\mathbf{W}^*)\|_{Z_{\eta/2,0}} \leq C \|\mathbf{W}^*\|_{Z'_{\eta,b_0}}.$$

Thus we have (35).

5.2 Stability of shooting solutions

To study the stability of the shooting solutions to (1) we can just as well study the stability of the solution \mathbf{W}^* to (31). Letting $\mathbf{W} = \mathbf{W}^* + \mathbf{V}$ and inserting this into (31) we find that \mathbf{V} satisfies

$$\mathbf{V}_t = \mathbf{A}\mathbf{V} + \mathbf{B}(0)\mathbf{V} + \mathbf{H}(\mathbf{V}) \tag{36}$$

with

$$\mathbf{V}(\mathbf{x}, -T) = \mathbf{V}_0(\mathbf{x}).$$

Here

$$\mathbf{H}(\mathbf{V}) := \mathbf{J}_{shoot}(\mathbf{W}^* + \mathbf{V}) - \mathbf{J}_{shoot}(\mathbf{W}^*)$$

(Since \mathbf{W}^* is defined for $t \leq 0$, we choose here to shift the initial data to $-T \leq 0$ instead of shifting \mathbf{W}^* forward in time.) The definition of \mathbf{J}_{shoot} gives:

$$\mathbf{H}(\mathbf{V}) = \mathbf{J}_{int}(\mathbf{W}^* + \mathbf{V}) - \mathbf{J}_{int}(\mathbf{W}^*) + \mathbf{J}_{nl}(\mathbf{W}^* + \mathbf{V}) - \mathbf{J}_{nl}(\mathbf{W}^*)$$

with $\mathbf{J}_{int} = (J_{1,int}, J_{2,int})$ and $\mathbf{J}_{nl} = (J_{1,nl}, J_{2,nl})$. A direct computation shows that

$$\begin{aligned} \mathbf{J}_{int}(\mathbf{W}^* + \mathbf{V}) - \mathbf{J}_{int}(\mathbf{W}^*) &= (\chi_1(\mathbf{x}, t) [F'(Q(\mathbf{y}) + Q(\mathbf{y}_2)) - F'(Q(\mathbf{y}))] V_1(\mathbf{y}), \\ & \quad \chi_2(\mathbf{x}, t) [F'(Q(\mathbf{y}_1) + Q(\mathbf{y})) - F'(Q(\mathbf{y}))] V_2(\mathbf{y})) \end{aligned}$$

Note that this term is linear in \mathbf{V} , unlike \mathbf{J}_{int} on its own, which contains an inhomogeneous term. This means that we can improve the similar estimate in Lemma 21 to:

$$\|\mathbf{J}_{int}(\mathbf{W}^* + \mathbf{V}) - \mathbf{J}_{int}(\mathbf{W}^*)\|_{L^5} \leq C e^{-\frac{b_0}{2}|\Delta\mathbf{r}_0|} \|\mathbf{V}\|_{L^5}.$$

Recalling that \mathbf{J}_{nl} consist of terms which are $O(|\mathbf{W}|^2)$, it is straightforward to conclude the pointwise estimate

$$|\mathbf{J}_{nl}(\mathbf{W}^* + \mathbf{V}) - \mathbf{J}_{nl}(\mathbf{W}^*)| \leq C \left(|\mathbf{W}^*| |\mathbf{V}| + |\mathbf{V}|^2 \right)$$

which implies

$$\|\mathbf{J}_{nl}(\mathbf{W}^* + \mathbf{V}) - \mathbf{J}_{nl}(\mathbf{W}^*)\|_{L^5} \leq C \left(\|\mathbf{W}^*\|_{L^\infty(\mathbf{R}^2)} \|\mathbf{V}\|_{L^5} + \|\mathbf{V}\|_{W^{8/5,5}}^2 \right).$$

The embedding (6) tells us that

$$\sup_{t \leq -T_3} \|\mathbf{W}^*\|_{L^\infty(\mathbf{R}^2)} \leq C e^{\eta T_3} \|\mathbf{W}^*\|_{Z'_{\eta,b}} \leq C e^{-\frac{b_0}{8}c|\Delta\mathbf{v}_0|T_3}.$$

Since \mathbf{B} is not small for general L^5 functions, and we wish to allow arbitrary initial data, we introduce, as before, the decomposition $\mathbf{V} = \mathbf{V}^{ic} + \mathbf{V}^{loc}$ where

$$\begin{aligned} \mathbf{V}_t^{ic} &= \mathbf{A}\mathbf{V}^{ic} + \mathbf{H}(\mathbf{V}) \\ \mathbf{V}_t^{loc} &= \mathbf{A}\mathbf{V}^{loc} + \mathbf{B}(0)\mathbf{V}. \end{aligned} \tag{37}$$

Given that \mathbf{A} has a triple eigenvalue, the above estimates will not allow us to conclude that \mathbf{V} is exponentially decaying. Note here that we have already specified $\dot{\mathbf{m}} = 0$ and so we cannot demand that $\mathbf{V} \in E^s \times E^s$ as we did when proving that stability of exit solutions. Instead of the Cauchy problem for (37), we proceed as in [19] and [21] and consider the boundary value problem:

$$\begin{aligned} \mathbf{P}^s \mathbf{V}^{ic}(\mathbf{x}, t = -T) &= \mathbf{V}^s, \quad \mathbf{P}^c \mathbf{V}^{ic}(\mathbf{x}, t = -T_3) = 0 \\ \mathbf{P}^s \mathbf{V}^{loc}(\mathbf{x}, t = -T) &= 0, \quad \mathbf{P}^c \mathbf{V}^{loc}(\mathbf{x}, t = -T_3) = 0. \end{aligned}$$

A solution of the boundary value problem will be a fixed point of the modified Duhamel integral function:

$$\begin{aligned} \Psi^{ic}(\mathbf{V}^{ic}, \mathbf{V}^{loc}) &:= e^{\mathbf{A}(t+T)} \mathbf{P}^s \mathbf{V}^s + \int_{-T}^t e^{\mathbf{A}(t-\tau)} \mathbf{P}^s \mathbf{H}(\mathbf{V}(\tau)) d\tau - \int_t^{-T_3} e^{\mathbf{A}(t-\tau)} \mathbf{P}^c \mathbf{H}(\mathbf{V}(\tau)) d\tau \\ \Psi^{loc}(\mathbf{V}^{ic}, \mathbf{V}^{loc}) &:= \int_{-T}^t e^{\mathbf{A}(t-\tau)} \mathbf{P}^s \mathbf{B}(0) (\Gamma^{ic}(\tau) + \mathbf{V}^{loc}(\tau)) d\tau - \int_t^{-T_3} e^{\mathbf{A}(t-\tau)} \mathbf{P}^c \mathbf{B}(0) (\Gamma^{ic}(\tau) + \mathbf{V}^{loc}(\tau)) d\tau. \end{aligned}$$

We compress notation: $X^b = X'_{\alpha/2,0}[T - T_3] \times X'_{\alpha/2,b_0}[T - T_3]$, $\mathbf{V}^b = (\mathbf{V}^{ic}, \mathbf{V}^{loc})$ and $\Psi^b = (\Psi^{ic}, \Psi^{loc})$. The third estimate in Theorem 15, along with Lemma 19 and those for \mathbf{H} above give the following estimates:

$$\begin{aligned} \|\Psi^b(\mathbf{V}^b)\|_{X^b} &\leq C^* \left(\|\mathbf{V}^s\|_{W^{8/5,5}} + e^{-\frac{b_0}{2}|\Delta\mathbf{r}_0|} \|\mathbf{V}^b\|_{X^b} + \|\mathbf{V}^b\|_{X^b}^2 \right) \\ \|\Psi^b(\mathbf{V}^b) - \Psi^b(\tilde{\mathbf{V}}^b)\|_{X^b} &\leq C^* \left(e^{-\frac{b_0}{2}|\Delta\mathbf{r}_0|} + \|\mathbf{V}^b\|_{X^b} + \|\tilde{\mathbf{V}}^b\|_{X^b} \right) \|\mathbf{V}^b - \tilde{\mathbf{V}}^b\|_{X^b} \end{aligned} \tag{38}$$

where C^* is a constant independent of T . We have in the above estimate taken T_3 sufficiently large so that $e^{-\frac{b_0}{8}c|\Delta\mathbf{v}_0|T_3} \leq e^{-\frac{b_0}{2}|\Delta\mathbf{r}_0|}$.

Let $\mathcal{K}_2 = 4C^*$ and $\delta_0 = \frac{1}{32(C^*)^2}$. If $\|\mathbf{V}^s\|_{W^{8/5,5}} = \delta \leq \delta_0$, and $e^{-\frac{b_0}{2}|\Delta\mathbf{r}_0|} \leq 1/\mathcal{K}_2$ then the last two estimates imply that Ψ^b is a contraction on the ball in X^b of radius $\mathcal{K}_2\delta$. And so there is a fixed point and the boundary value problem (37) has a solution

$$\|\mathbf{V}^b\|_{X^b} \leq \mathcal{K}_2 \|\mathbf{V}^s\|_{W^{8/5,5}}$$

where the constant \mathcal{K}_2 is independent of T .

Let

$$\mathcal{T}^c(\mathbf{V}^s; \mathbf{n}) = \mathbf{P}^c(\mathbf{V}^{ic}(t = -T) + \mathbf{V}^{loc}(t = -T))$$

where $(\mathbf{V}^{ic}, \mathbf{V}^{loc})$ is the solution of the boundary value problem. Here

$$\mathbf{n} = (\mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20}).$$

If we were in a situation where $\mathbf{V}_0 = \mathbf{V}^s + \mathcal{T}^c(\mathbf{V}^s, \mathbf{n})$ for some function $\mathbf{V}^s \in E^s \times E^s$, then the solution of the initial value problem for (37) would decay exponentially quickly and we would be done.

This will not generally be the case, but it turns out that this intuition is enough to complete the proof. Set

$$\mathbf{Q}(\mathbf{x}; \mathbf{n}) = (Q(\mathbf{y}_1(\mathbf{n})), Q(\mathbf{y}_2(\mathbf{n})))$$

and (with a small abuse of notation)

$$\mathbf{W}^*(\mathbf{x}; \mathbf{n}) = (W_1^*(\mathbf{y}_1, -T), W_2^*(\mathbf{y}_2, -T))$$

where $\mathbf{y}_j(\mathbf{n}) = \mathcal{R}[\theta_{j0}](\mathbf{x} - \mathbf{r}_{j0} + \mathbf{v}(\theta_{j0})cT)$. Notice that if we could find $\mathbf{V}^s \in E^s \times E^s$ and \mathbf{n}_1 so that

$$\mathbf{Q}(\mathbf{x}; \mathbf{n}) + \mathbf{W}^*(\mathbf{x}; \mathbf{n}) + \mathbf{V}_0 = \mathbf{Q}(\mathbf{x}; \mathbf{n}_1) + \mathbf{W}^*(\mathbf{x}; \mathbf{n}_1) + \mathbf{V}^s + \mathcal{T}^c(\mathbf{V}^s; \mathbf{n}_1) \quad (39)$$

then we would be done. That is to say, if the slight adjustment \mathbf{n}_1 of \mathbf{n} can be found so that we have the above, then our initial conditions would be such that the boundary value problem and the initial value problem coincide and we have the exponential decay. (Note that in the above we make sure that the components of \mathbf{V}_0 , \mathbf{V}^s and $\mathcal{T}^c(\mathbf{V}^s; \mathbf{n}_1)$ are evaluated at the appropriate $\mathbf{y}_j(\mathbf{n})$.)

The proof of the claim follows from the application of the implicit function theorem to the map

$$P(\mathbf{V}^s, \mathbf{n}_1; \mathbf{V}^0) = \mathbf{Q}(\mathbf{x}; \mathbf{n}_1) + \mathbf{W}^*(\mathbf{x}; \mathbf{n}_1) + \mathbf{V}^s + \mathcal{T}^c(\mathbf{V}^s; \mathbf{n}_1) - \mathbf{Q}(\mathbf{x}; \mathbf{n}) - \mathbf{W}^*(\mathbf{x}; \mathbf{n}) - \mathbf{V}_0.$$

Clearly $P(0, 0; \mathbf{n}) = 0$. Notice $D_{\mathbf{V}^s} \mathbf{P}^s P(0, 0; \mathbf{n}) = id_{E^s \times E^s}$ and $D_{\mathbf{V}^s} \mathbf{P}^c P(0, 0; \mathbf{n}) = D_{\mathbf{V}^s} \mathcal{T}^c(0; \mathbf{n})$ since $\mathbf{V}^s \in E^s \times E^s$ and $\mathcal{T}^c \in E^c \times E^c$. It is the derivatives of Q with respect to the parameters \mathbf{r}_{j0} and θ_{j0} which give the center eigenspace and so

$$D_{\mathbf{n}_1} \mathbf{P}^c \mathbf{Q}(\mathbf{x}, \mathbf{n}_1) = id_6$$

where id_6 is the six-by-six identity matrix and

$$D_{\mathbf{n}_1} \mathbf{P}^s \mathbf{Q}(\mathbf{x}, \mathbf{n}_1) = 0.$$

From (35) we conclude that $\|D_{\mathbf{n}_1} \mathbf{W}^*(\mathbf{x}, \mathbf{n}_1)\|_{W^{8/5,5}} \leq Ce^{-\frac{b_0}{16}c|\Delta\mathbf{v}_0|T}$. Finally, $\mathcal{T}^c(0; \mathbf{n}) = 0$ for all choices \mathbf{n} and so $D_{\mathbf{n}} \mathcal{T}^c(0; \mathbf{n}) = 0$. Decomposing P into $(\mathbf{P}^s P, \mathbf{P}^c P)$ and arranging all the derivatives computed above accordingly, we see

$$D_{(\mathbf{V}^s, \mathbf{n})} P(0, \mathbf{n}; 0) = \begin{bmatrix} id_{E^s \times E^s} & D_{\mathbf{V}^s} \mathcal{T}^c(0; \mathbf{n}) \\ 0 & id_6 \end{bmatrix} + O(e^{-\frac{b_0}{8}c|\Delta\mathbf{v}_0|T}).$$

This is invertible if T is sufficiently large, which it is. This completes the proof.

We conclude this section with the proof of Corollary 10:

Proof. Suppose that $\phi_j(\mathbf{x}, t; \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20})$ for $j = 1, 2$ are two different solutions of (1) meeting the conclusions of Theorem 7. Fix $T > T_3$, let

$$h = \|\phi_1(\cdot, -T, \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20}) - \phi_2(\cdot, -T, \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20})\|_{W^{8/5,5}} \neq 0$$

We know that there is a $T_2 > T$ so that for $j = 1, 2$ and any $\gamma > 0$

$$\left\| \phi_j(\cdot, -T_2, \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20}) - \sum_{j=1}^2 Q(\mathcal{R}[-\theta_{j0}] (\cdot - \mathbf{r}_j^*(-T_2))) \right\|_{W^{8/5,5}} \leq \gamma.$$

which then implies by the triangle inequality:

$$\|\phi_1(\cdot, -T, \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20}) - \phi_2(\cdot, -T, \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20})\|_{W^{8/5,5}} \leq 2\gamma.$$

Applying Theorem 9 gives

$$\|\phi_1(\cdot, -T_2, \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20}) - \phi_2(\cdot, -T_2, \mathbf{r}_{10}, \mathbf{r}_{20}, \theta_{10}, \theta_{20})\|_{W^{8/5,5}} \leq 2\mathcal{K}_3\gamma e^{\eta(T_2-T)} \leq 2\mathcal{K}_3\gamma.$$

Since γ is free to choose, we can take it to be $h/20\mathcal{K}_3$. This is a contradiction. \square

6 Appendix: Local existence of solutions for (20)

Let

$$\mathbf{S} = (\mathbf{W}^{ic}, \mathbf{W}^{loc}, \mathbf{m}),$$

$$Y[T_1] := \in X'_{a,0}[T_1] \times X'_{a,b}[T_1] \times C^1[0, T_1]$$

and

$$Y_0 := (W^{4/3,3} \cap (E^s)^2) \times (W_b^{4/3,3} \cap (E^s)^2) \times \mathbf{R}^3.$$

Theorem 27. *There exists $C > 0$ and $\delta_2 > 0$ so that for any $a > 0$, $0 \leq b \leq b_0$ and*

$$\|\mathbf{S}_0\|_{Y_0} \leq \delta_2$$

there exists $T_1 > 0$ and a unique function and $\mathbf{S} \in Y[T_1]$ so that

$$\|\mathbf{S}\|_{Y[T_1]} \leq C$$

and \mathbf{S} solves (20) for $t \in [0, T_1]$ a.e. and $\mathbf{S}(0) = \mathbf{S}_0$.

Proof. Let $\Gamma(\mathbf{S}) := (\Gamma^{ic}(\mathbf{S}), \Gamma^{loc}(\mathbf{S}), \Gamma^m(\mathbf{S}))$ be the map defined by the following, which is formally equivalent to the the right hand side of (21):

$$\begin{aligned} \Gamma^{ic} &= e^{\mathbf{A}t} \mathbf{W}_0^{ic} + \int_0^t e^{\mathbf{A}(t-\tau)} \mathbf{P}^s \mathbf{J}_s(\mathbf{m}(\tau), \mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau)) d\tau \\ \Gamma^{loc} &= e^{\mathbf{A}t} \mathbf{W}_0^{loc} + \int_0^t e^{\mathbf{A}(t-\tau)} \mathbf{P}^s \mathbf{B}(\mathbf{m}(\tau)) (\mathbf{W}^{ic}(\tau) + \mathbf{W}^{loc}(\tau)) d\tau \\ \Gamma^m &= \mathbf{m}_0 - \int_0^t \{ \mathbf{R}(\mathbf{m}(\tau)) \mathbf{P}^c (\mathbf{B}(\mathbf{m}(\tau)) (\Gamma^{ic}(\tau) + \Gamma^{loc}(\tau)) \\ &\quad + \mathbf{J}_{cc}(\mathbf{m}(\tau), \Gamma^{ic}(\tau) + \Gamma^{loc}(\tau))) + \mathbf{R}[\mathbf{m}(\tau)] \mathbf{K}_{cc}(\mathbf{m}(\tau), \Gamma^{ic}(\tau) + \Gamma^{loc}(\tau)) \} d\tau. \end{aligned} \tag{40}$$

We will show that that there exists $T_1 > 0$, $K^{**} > 0$ and $\delta_2 > 0$ so that $\|\mathbf{S}_0\|_{Y_0} \leq \delta_2$ and $\|\mathbf{S}\|_Y \leq K^{**}$ imply

$$\|\Gamma(\mathbf{S})\|_Y \leq K^{**} \tag{41}$$

and

$$\left\| \Gamma(\mathbf{S}) - \Gamma(\tilde{\mathbf{S}}) \right\|_Y \leq \frac{1}{2} \left\| \mathbf{S} - \tilde{\mathbf{S}} \right\|_Y \quad (42)$$

which by the contraction mapping theorem implies the result.

First we estimate Γ^{ic} . Theorem 15 gives:

$$\left\| \Gamma^{ic}(\mathbf{S}) \right\|_{X_{a,0}[T]} \leq C \left(\left\| \mathbf{S}_0 \right\|_{Y_0} + \left\| \mathbf{J} \right\|_{X_{a,0}[T]} \right).$$

Lemma (20)-(23) imply

$$\begin{aligned} \left\| \mathbf{J} \right\|_{X_{a,0}[T]} \leq C & \left(\left(\int_0^T e^{-3b\mu(t)/2} dt \right)^{1/3} + \left\| \mathbf{W}^{ic} \right\|_{L^5([0,T]; W^{4/3,3})} + \left\| \mathbf{W}^{loc} \right\|_{L^5([0,T]; W_b^{4/3,3})} \right. \\ & \left. + \left\| \mathbf{m} \right\|_{C^1[0,T]}^2 + \left\| \mathbf{W}^{ic} \right\|_{X'_{a,0}[T]}^2 + \left\| \mathbf{W}^{loc} \right\|_{X'_{a,b}[T]}^2 \right) \quad (43) \end{aligned}$$

To estimate the linear terms above we use (6) and Hölder's inequality:

$$\begin{aligned} \left\| \mathbf{F} \right\|_{L^5([0,T], W_b^{4/3,3})} &= \left(\int_0^T \left\| \mathbf{F}(\tau) \right\|_{W_b^{4/3,3}}^3 d\tau \right)^{1/3} \\ &\leq \left\| \mathbf{F} \right\|_{L^\infty([0,T], W_b^{4/3,3})}^{2/3} \left(\int_0^T \left\| \mathbf{F}(\tau) \right\|_{W_b^{4/3,3}} d\tau \right)^{1/3} \\ &\leq C \left\| \mathbf{F} \right\|_{X'_{a,b}}^{2/3} \left(\int_0^T \left\| \mathbf{F}(\tau) \right\|_{W_b^{4/3,3}} d\tau \right)^{1/3} \\ &\leq CT^{2/9} \left\| \mathbf{F} \right\|_{X'_{a,b}}. \end{aligned}$$

Thus

$$\left\| \Gamma^{ic}(\mathbf{S}) \right\|_{X'_{a,0}} \leq C \left\{ \left\| \mathbf{S}_0 \right\|_{Y_0} + T^{2/9} (1 + \left\| \mathbf{S} \right\|_Y) + \left\| \mathbf{S} \right\|_Y^2 \right\}.$$

In exactly the same fashion we can show

$$\left\| \Gamma^{loc}(\mathbf{S}) \right\|_{X'_{a,0}} \leq C \left\{ \left\| \mathbf{S}_0 \right\|_{Y_0} + T^{2/9} \left\| \mathbf{S} \right\|_Y \right\}$$

and

$$\left\| \Gamma^{\mathbf{m}} \right\|_{C^1[0,T]} \leq C \left\{ \left\| \mathbf{S}_0 \right\|_{Y_0} + T^{2/9} (1 + \left\| \mathbf{S} \right\|_Y) + \left\| \mathbf{S} \right\|_Y^2 + \left\| \mathbf{S} \right\|_Y^3 \right\}.$$

Therefore

$$\left\| \Gamma(\mathbf{S}) \right\|_Y \leq C \left\| \mathbf{S}_0 \right\|_{Y_0} + CT^{2/9} (1 + \left\| \mathbf{S} \right\|_Y) + C \left\| \mathbf{S} \right\|_Y^2 + C \left\| \mathbf{S} \right\|_Y^3$$

which in turn implies (41) for T and $\left\| \mathbf{S}_0 \right\|_{Y_0}$ sufficiently small.

Proving (42) is largely similar, though care must be taken with $\mathbf{B}(\mathbf{m})\mathbf{W}$ (as well as some terms in $J_{j,int}$) because it is not immediately obvious that this term is Lipschitz. Of particular concern is that fact that these operators involve spatial translations and thus are nonlocal.

The most difficult term is Γ^{loc}

$$\begin{aligned} \Gamma^{loc}(\mathbf{S}) - \Gamma^{loc}(\tilde{\mathbf{S}}) &= \int_0^t e^{\mathbf{A}(t-\tau)} \mathbf{P}^s \mathbf{B}(\mathbf{m}(\tau)) \left(\mathbf{W}(\tau) - \tilde{\mathbf{W}}(\tau) \right) d\tau \\ &\quad - \int_0^t e^{\mathbf{A}(t-\tau)} \mathbf{P}^s \left(\mathbf{B}(\mathbf{m}(\tau)) - \mathbf{B}(\tilde{\mathbf{m}}(\tau)) \right) \mathbf{W}(\tau) d\tau. \end{aligned}$$

Here we use the shorthand $\mathbf{W} = \mathbf{W}^{ic} + \mathbf{W}^{loc}$. The first term we can estimate by $CT^{2/9} \|\mathbf{S} - \tilde{\mathbf{S}}\|_Y$ using the same strategy used for (41).

We claim that:

$$\|(\mathbf{B}(\mathbf{m}(t)) - \mathbf{B}(\tilde{\mathbf{m}}(t))) \mathbf{W}(t)\|_{L_b^5} \leq C(1+t) \|\mathbf{W}\|_{W^{8/5,5}} |\mathbf{m} - \tilde{\mathbf{m}}| \quad (44)$$

From the definition of \mathbf{B} and the operators B_j we see that we must estimate terms of the form

$$\begin{aligned} & |\chi_1(\mathbf{x}, t) [F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))] W_2(\mathbf{y}_2, t) \\ & - \chi_1(\tilde{\mathbf{x}}, t) [F'(Q(\tilde{\mathbf{z}}_1) + Q(\tilde{\mathbf{z}}_2)) - F'(Q(\tilde{\mathbf{z}}_2))] W_2(\tilde{\mathbf{y}}_2, t)| \\ \leq & |\chi_1(\mathbf{x}, t) [F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))] (W_2(\mathbf{y}_2, t) - W_2(\tilde{\mathbf{y}}_2, t))| \\ & + |\chi_1(\mathbf{x}, t) [F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))] - \chi_1(\tilde{\mathbf{x}}, t) [F'(Q(\tilde{\mathbf{z}}_1) + Q(\tilde{\mathbf{z}}_2)) - F'(Q(\tilde{\mathbf{z}}_2))]| |W_2(\tilde{\mathbf{y}}_2)| \end{aligned} \quad (45)$$

$W_2(t)$ is in $W^{8/5,5}$ and thus is C^1 . So:

$$\begin{aligned} & |\chi_1(\mathbf{x}, t) [F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))] (W_2(\mathbf{y}_2, t) - W_2(\tilde{\mathbf{y}}_2, t))| \\ & \leq C |\chi_1(\mathbf{x}, t) [F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))]| |\mathbf{y}_2 - \tilde{\mathbf{y}}_2| \|W_2\|_{W^{8/5,5}} \end{aligned}$$

From Remark 18 (and noting that \mathbf{m} and $\tilde{\mathbf{m}}$ have identical initial data) we have

$$\mathbf{y}_2 - \tilde{\mathbf{y}}_2 = \mathcal{R}[-\theta_{20}] (\mathbf{r}_2 - \tilde{\mathbf{r}}_2 - \mathbf{r}_1 + \tilde{\mathbf{r}}_1).$$

Then

$$\begin{aligned} |\mathbf{y}_2 - \tilde{\mathbf{y}}_2| & \leq \sum_j |\mathbf{r}_j - \tilde{\mathbf{r}}_j| \\ & \leq \sum_j |\mathbf{p}_j - \tilde{\mathbf{p}}_j| + c \int_0^t \left| \mathcal{R}[\theta_{20} + \phi_j(s)] - \mathcal{R}[\theta_{20} + \tilde{\phi}_j(s)] \right| ds \\ & \leq C(1+t) |\mathbf{m} - \tilde{\mathbf{m}}|. \end{aligned}$$

Additionally, we know from the proof of Lemma 19 that:

$$\|[F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))]\| \leq Ce^{-\beta|\mathbf{y}|}$$

which is in L_b^5 . Therefore

$$\begin{aligned} & \|\chi_1(\mathbf{x}, t) [F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))] (W_2(\mathbf{y}_2, t) - W_2(\tilde{\mathbf{y}}_2, t))\|_{L_b^5} \\ & \leq C(1+t) \|\mathbf{W}\|_{W^{8/5,5}} |\mathbf{m} - \tilde{\mathbf{m}}|. \end{aligned}$$

The remaining term in (45) is handled in a completely similar way, even though it is quite awful looking. Let

$$f(\mathbf{y}, t; \mathbf{m}) := e^{b|\mathbf{y}|} \chi_1(\mathbf{x}, t) [F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))].$$

This function is Lipschitz in its arguments (since χ_1 , Q and F' are all Lipschitz) and also in L^∞ . Thus

$$\|f(\mathbf{y}, t; \mathbf{m}) - f(\mathbf{y}, t; \tilde{\mathbf{m}})\|_{L^\infty} \leq C(1+t) |\mathbf{m} - \tilde{\mathbf{m}}|.$$

From this and Remark 18 we can conclude

$$\begin{aligned} & \|\chi_1(\mathbf{x}, t) [F'(Q(\mathbf{z}_1) + Q(\mathbf{z}_2)) - F'(Q(\mathbf{z}_2))] \\ & - \chi_1(\tilde{\mathbf{x}}, t) [F'(Q(\tilde{\mathbf{z}}_1) + Q(\tilde{\mathbf{z}}_2)) - F'(Q(\tilde{\mathbf{z}}_2))] W_2(\tilde{\mathbf{y}}_2)\|_{L_b^5} \\ & \leq C(1+t) \|\mathbf{W}\|_{W^{8/5,5}} |\mathbf{m} - \tilde{\mathbf{m}}|. \end{aligned}$$

Thus we have established the claim (44) which in turn implies

$$\left\| \Gamma^{loc}(\mathbf{S}) - \Gamma^{loc}(\tilde{\mathbf{S}}) \right\|_Y \leq C \left(T^{2/9} + (1+T) \left(\|\mathbf{S}\|_Y + \|\tilde{\mathbf{S}}\|_Y \right) \right) \|\mathbf{S} - \tilde{\mathbf{S}}\|_Y$$

Arguments parallel to this can be used to establish similar estimates for Γ^{ic} and Γ^m which then implies (42). We omit the details. \square

References

- [1] J. C. Alexander and C. K. R. T. Jones. Existence and stability of asymptotically oscillatory double pulses. *J. Reine Angew. Math.*, 446:49–79, 1994.
- [2] H. Amann. *Linear and quasilinear parabolic problems. Vol. I*, volume 89 of *Monographs in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1995. Abstract linear theory.
- [3] M. Bär, N. Gottschalk, M. Eiswirth, and G. Ertl. Spiral waves in a surface reaction: model calculations. *J. Chem. Phys.*, 100(2):1202–1214, 1994.
- [4] W.-J. Beyn, S. Selle, and V. Thümler. Freezing multipulses and multifronts. *SIAM J. Appl. Dyn. Syst.*, 7(2):577–608, 2008.
- [5] X. Chen and J.-S. Guo. Existence and uniqueness of entire solutions for a reaction-diffusion equation. *J. Differential Equations*, 212(1):62–84, 2005.
- [6] S.-I. Ei. The motion of weakly interacting pulses in reaction-diffusion systems. *J. Dynam. Differential Equations*, 14(1):85–137, 2002.
- [7] J. W. Evans, N. Fenichel, and J. A. Feroe. Double impulse solutions in nerve axon equations. *SIAM J. Appl. Math.*, 42(2):219–234, 1982.
- [8] J. A. Feroe. Existence and stability of multiple impulse solutions of a nerve equation. *SIAM J. Appl. Math.*, 42(2):235–246, 1982.
- [9] P. C. Fife. Long time behavior of solutions of bistable nonlinear diffusion equations. *Arch. Rational Mech. Anal.*, 70(1):31–46, 1979.
- [10] P. C. Fife and J. B. McLeod. The approach of solutions of nonlinear diffusion equations to travelling front solutions. *Arch. Ration. Mech. Anal.*, 65(4):335–361, 1977.
- [11] J.-S. Guo and Y. Morita. Entire solutions of reaction-diffusion equations and an application to discrete diffusive equations. *Discrete Contin. Dyn. Syst.*, 12(2):193–212, 2005.
- [12] S. P. Hastings. Single and multiple pulse waves for the FitzHugh-Nagumo equations. *SIAM J. Appl. Math.*, 42(2):247–260, 1982.
- [13] D. Henry. *Geometric theory of semilinear parabolic equations*, volume 840 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1981.
- [14] Y. Morita and H. Ninomiya. Entire solutions with merging fronts to reaction-diffusion equations. *J. Dynam. Differential Equations*, 18(4):841–861, 2006.
- [15] Y. Nishiura, T. Teramoto, and K.-I. Ueda. Scattering of traveling spots in dissipative systems. *Chaos*, 15(4):047509, 10, 2005.
- [16] A. Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1983.
- [17] H.-G. Purwins, H. U. Bödeker, and A. W. Liehr. Dissipative solitons in reaction-diffusion systems. In *Dissipative solitons*, volume 661 of *Lecture Notes in Phys.*, pages 267–308. Springer, Berlin, 2005.

- [18] B. Sandstede. Stability of multiple-pulse solutions. *Trans. Amer. Math. Soc.*, 350(2):429–472, 1998.
- [19] A. Scheel and J. D. Wright. Colliding dissipative pulses—the shooting manifold. *J. Differential Equations*, 245(1):59–79, 2008.
- [20] L. Tartar. *An introduction to Sobolev spaces and interpolation spaces*, volume 3 of *Lecture Notes of the Unione Matematica Italiana*. Springer, Berlin, 2007.
- [21] J. D. Wright. Separating dissipative pulses: the exit manifold. *J. Dynam. Differential Equations*, 21(2):315–328, 2009.
- [22] H. Yagisita. Backward global solutions characterizing annihilation dynamics of travelling fronts. *Publ. Res. Inst. Math. Sci.*, 39(1):117–164, 2003.
- [23] S. Zelik and A. Mielke. Multi-pulse evolution and space-time chaos in dissipative systems. *preprint*, 2006.