



Average Number of Distinct Part Sizes in a Random Carlitz Composition

WILLIAM M. Y. GOH AND PAWEŁ HITCZENKO

A composition of an integer n is called Carlitz if adjacent parts are different. Several characteristics of random Carlitz compositions have been studied recently by Knopfmacher and Prodinger. We will complement their work by establishing asymptotics of the average number of distinct part sizes in a random Carlitz composition.

© 2002 Published by Elsevier Science Ltd.

1. INTRODUCTION

In this paper we obtain precise asymptotics, as $n \rightarrow \infty$, for the expected number of distinct part sizes in a random Carlitz composition of an integer n . Let us recall that a tuple $(\gamma_1, \dots, \gamma_k)$ is a *composition* of an integer n if the γ_j 's are positive integers, called parts, such that $\sum_j \gamma_j = n$. The number k is the number of parts and the values of γ_j 's are called part sizes. There are 2^{n-1} different compositions of n . A composition is called *Carlitz* if the adjacent parts are different, i.e., if $\gamma_j \neq \gamma_{j+1}$ for $j = 1, \dots, k-1$. For example, out of 16 compositions of the integer 5, seven are Carlitz, namely (5), (4, 1), (1, 4), (3, 2), (2, 3), (1, 3, 1), and (2, 1, 2). For other values, see [8, Sequence A003242]. We denote the set of all Carlitz compositions of n by Ω_n . Carlitz compositions have been introduced by Carlitz [1] who found the generating function for the total number of them. They have been subsequently studied by Knopfmacher and Prodinger [5] (see also [6]). These authors found the asymptotics of the total number of Carlitz compositions. They also studied several parameters (like the number of parts, the size of the largest part, among other things) for *random* Carlitz compositions. ‘Random Carlitz composition’ means a composition chosen according to the uniform probability measure on Ω_n . This measure will be denoted by \mathbf{P} and \mathbf{E} will denote integration with respect to \mathbf{P} . In this setting, various parameters of Carlitz compositions become random variables and their probabilistic properties are to be studied. For example Knopfmacher and Prodinger found, among other things, the exact asymptotic behavior of the expected number of parts and the expected size of the largest part. One question that they left open concerned the expected value of the number of *distinct* part sizes, D_n , and the purpose of this note is to answer their question. In order to state the result we need to introduce some more notation and we will try to closely follow the notation of Knopfmacher and Prodinger. First, the number of distinct part sizes is defined formally as follows: if $(\gamma_1, \dots, \gamma_k)$ is in Ω_n then

$$D_n = 1 + \sum_{i=2}^k I_{\{\gamma_i \neq \gamma_j, j=1, \dots, i-1\}},$$

where I_A denotes, as usual, the indicator function of the set A . Secondly, we introduce a function σ of a complex variable defined by

$$\sigma(z) = \sum_{j=1}^{\infty} (-1)^{j-1} \frac{z^j}{1-z^j}.$$

The equation $\sigma(z) = 1$ has the unique real solution $\rho = 0.571349\dots$ on the interval $[0, 1]$. (The relevance of this is that, as was shown by Carlitz, the generating function of Carlitz compositions is equal to $1/(1-\sigma(z))$.)

THEOREM 1. *With the above notation, and letting $\{\cdot\}$ denote the fractional part, as $n \rightarrow \infty$, we have*

$$\mathbf{E}D_n = \frac{\ln(n/\sigma'(\rho))}{\ln(1/\rho)} + \frac{1}{2} - \frac{\gamma}{\ln(1/\rho)} + h_0(\{\rho \ln(n/\sigma'(\rho))\}) + o(1),$$

where γ is Euler's constant, h_0 is a mean zero function of period 1 whose Fourier coefficients are given by

$$c_\ell = \frac{1}{\ln(1/\rho)} \Gamma\left(-\frac{2\pi i \ell}{\ln(1/\rho)}\right), \quad \ell \neq 0.$$

Approximating all the constants and using the fact that the gamma function decays very fast along the imaginary axis, it can be seen, in particular, that

$$\mathbf{E}D_n = C_1 \ln n - C_2 + h_0(\{\rho \ln(n/\sigma'(\rho))\}) + o(1),$$

where $C_1 = 1.786495\dots$, $C_2 = 2.932545\dots$, and the amplitude of h_0 is bounded by $0.5882304\dots \times 10^{-7}$.

The approach is as in [5] via generating functions. We let I_j denote the set of those Carlitz compositions that contain at least one part of size ' j '. Using, without any risk of confusion, the same notation for a set and its indicator we have

$$D_n = \sum_{j=1}^n I_j,$$

and therefore

$$\mathbf{E}D_n = \sum_{j=1}^n \mathbf{P}(I_j) = \sum_{j=1}^n (1 - \mathbf{P}(I_j^c)), \quad (1)$$

where A^c denotes the complement of a set A . Denoting by a_n and $a_{n,j}$ the number of all Carlitz compositions of n and the number of those Carlitz compositions of n that *do not* use size j we have

$$\mathbf{P}(I_j^c) = \frac{a_{n,j}}{a_n}.$$

The sequence (a_n) was studied in Knopfmacher and Prodinger [5], so we need only to study the numbers $a_{n,j}$. In order to do that we will build their generating function. The construction follows the ideas of [5], but since that paper is short on some details and may be a bit difficult to read for a new adept, we will provide a fairly detailed argument.

We would like to mention that the method of our paper does not appear to be strong enough to yield information about the limiting distribution of the variable D_n . A bivariate generating function for D_n would be extremely welcome. But, we were unable to get it. Although it is known (cf. [6]) that the total number of parts in Carlitz composition satisfies the central limit theorem, we think that it is unlikely that D_n will have the same property. It would be very interesting and desirable to find the limiting distribution of D_n .

2. GENERATING FUNCTION

In this section we will prove the following statement

PROPOSITION 2. *Let $C_j(z)$ be the generating function of the sequence $\{a_{n,j}, n \geq 0\}$. Then,*

$$C_j(z) = \frac{1}{1 - \sigma_j(z)},$$

where

$$\sigma_j(z) = \sum_{\ell \geq 1} (-1)^{\ell-1} \left(\frac{z^\ell}{1-z^\ell} - z^{\ell j} \right) = \sigma(z) - \frac{z^j}{1-z^j}. \tag{2}$$

PROOF. Let $a_j^{(k)}(n, m)$ denote the number of Carlitz compositions of n with the properties:

- they have exactly k parts
- they do not contain a part of size j
- the last part is of size m .

Then for $k \geq 1$ and $m \neq j$ we have

$$a_j^{(k+1)}(n, m) = a_j^{(k)}(n - m) - a_j^{(k)}(n - m, m),$$

where $a_j^{(k)}(\ell)$ is the number of Carlitz compositions of ℓ into k parts, none of them equal to j . Since j is fixed throughout the argument, for the ease of notation we suppress the subscript j throughout the argument. We additionally require that $a^{(k)}(\ell)$ and $a^{(k)}(\ell, m)$ vanish whenever $\ell \leq 0$. Let

$$f_k(z, u) = \sum_{m \geq 1, n \geq 0} a^{(k)}(n, m) z^n u^m.$$

Since the compositions enumerated in $a^{(k)}(n, m)$ do not contain a part of size j , we have $a^{(k)}(n, j) = 0$. Hence

$$\begin{aligned} f_{k+1}(z, u) &= \sum_{n \geq 0} \sum_{m \geq 1} a^{(k+1)}(n, m) z^n u^m \\ &= \sum_{n \geq 0} \sum_{\substack{m \geq 1 \\ m \neq j}} a^{(k+1)}(n, m) z^n u^m + \sum_{n \geq 0} a^{(k+1)}(n, j) z^n u^j \\ &= \sum_{n \geq 0} \sum_{\substack{m \geq 1 \\ m \neq j}} (a^{(k)}(n - m) - a^{(k)}(n - m, m)) z^n u^m \\ &= \sum_{n \geq 0} \sum_{\substack{m \geq 1 \\ m \neq j}} a^{(k)}(n - m) z^n u^m \\ &\quad - \sum_{n \geq 0} \sum_{\substack{m \geq 1 \\ m \neq j}} a^{(k)}(n - m, m) z^n u^m. \end{aligned}$$

The first sum above is equal to

$$\begin{aligned} &\sum_{n \geq 0, m \geq 1} a^{(k)}(n - m) z^n u^m - \sum_{n \geq 0} a^{(k)}(n - j) z^n u^j \\ &= \sum_{\ell \geq 0, m \geq 1} a^{(k)}(\ell) (zu)^m z^\ell - \sum_{m \geq 1} a^{(k)}(m) z^m (zu)^j \\ &= f_k(z, 1) \frac{zu}{1-zu} - f_k(z, 1) (zu)^j \end{aligned}$$

where, in the second step we changed the summation indices $n - m = \ell$ and $n - j = m$, respectively. By the same argument

$$\sum_{m \geq 1, \ell \geq 0} a^{(k)}(\ell, m) z^\ell (zu)^m = f_k(z, zu).$$

Thus, for $k \geq 1$ we have

$$f_{k+1}(z, u) = f_k(z, 1) \left(\frac{zu}{1-zu} - (zu)^j \right) - f_k(z, zu).$$

Letting $f_0(z, u) = 1$ the last line can be rewritten as

$$f_{k+1}(z, u) = f_k(z, 1) \left(\frac{zu}{1-zu} - (zu)^j \right) - f_k(z, zu) + \delta_{k,0},$$

for $k \geq 0$. Introducing the function

$$F(z, u) = \sum_{k \geq 1} f_k(z, u)$$

and summing over $k \geq 0$ we obtain

$$F(z, u) = F(z, 1) \left(\frac{zu}{1-zu} - (zu)^j \right) + \left(\frac{zu}{1-zu} - (zu)^j \right) - F(z, zu).$$

This equation can be iterated to yield

$$\begin{aligned} F(z, u) &= F(z, 1) \left(\sum_{\ell \geq 1} (-1)^{\ell-1} \left(\frac{z^\ell u}{1-z^\ell u} - (z^\ell u)^j \right) \right) \\ &\quad + \sum_{\ell \geq 1} (-1)^\ell \left(\frac{z^\ell u}{1-z^\ell u} - (z^\ell u)^j \right). \end{aligned}$$

Hence, for $u = 1$ we get

$$F(z, 1) = F(z, 1)\sigma_j(z) + \sigma_j(z),$$

where $\sigma_j(z)$ is defined by (2). Finally, letting

$$C_j(z) = 1 + F(z, 1),$$

we see that

$$C_j(z) - 1 = (C_j(z) - 1)\sigma_j(z) + \sigma_j(z),$$

which, since

$$F(z, 1) = \sum_{k \geq 1} f_k(z, 1) = \sum_{k \geq 1} \sum_{\substack{n \geq 0 \\ m \geq 1}} a^{(k)}(n, m) z^n = \sum_{n \geq 0} a_{n,j} z^n,$$

means that

$$C_j(z) = \frac{1}{1 - \sigma_j(z)},$$

is the generating function of the sequence $(a_{n,j})$. □

3. SINGULARITIES OF THE GENERATING FUNCTION

A starting point of our analysis is the fact that the generating function of Carlitz compositions has the unique singularity in the disc $\{z : |z| \leq 0.663\}$. This singularity is the unique real root, ρ , of the equation $\sigma(z) = 1$ on $[0, 1]$. The numerical approximation of that root is $\rho = 0.571349\dots$. Since $\sigma(z)$ and $\sigma_j(z) = \sigma(z) - z^j/(1-z^j)$ do not differ by too much, functions σ_j will have the same feature, at least for j 's sufficiently large. In fact, on the disc $\{z : |z| \leq 0.663\}$ the functions satisfy the following: there exists $\delta > 0$ such that for every $j \geq 6$ the equation $\sigma_j(z) = 1$ has the unique real simple root on $[0, 1]$. Furthermore, these roots, which will be denoted by ρ_j , have the following properties

1. $\forall j \geq 6, 0 < \rho_j \leq \rho + \delta,$
2. $\forall j \geq 6$ all roots ξ of $\sigma_j(z) = 1$ other than ρ_j satisfy $|\xi| \geq \rho + 2\delta,$
3. ρ_j are strictly decreasing for $j \geq 6$ and $\rho_j \rightarrow \rho$ as $j \rightarrow \infty.$

A justification as well as a discussion of the remaining case $1 \leq j \leq 5$ is postponed until the appendix. We will need asymptotics of ρ_j and we will use the ‘bootstrapping method’. Rewriting

$$\sigma_j(z) = 1$$

as

$$\sigma(z) = \frac{1}{1 - z^j}, \tag{3}$$

letting $\rho_j = \rho + \varepsilon_j,$ where $\varepsilon_j = o(1)$ and substituting the latter expression for ρ_j into (3) we get

$$\sigma(\rho + \varepsilon_j) = \frac{1}{1 - (\rho + \varepsilon_j)^j} = \frac{1}{1 - \rho^j(1 + \varepsilon_j/\rho)^j}.$$

Using Taylor’s expansion on both sides we see that

$$\sigma(\rho) + \sigma'(\rho)\varepsilon_j + O(\varepsilon_j^2) = 1 + \rho^j \left(1 + \frac{\varepsilon_j}{\rho}\right)^j + O(\rho^2 j).$$

Since $\sigma(\rho) = 1$ we obtain that $\varepsilon_j = \rho^j/\sigma'(\rho) + o(\rho^j).$ Hence

$$\rho_j = \rho + \frac{\rho^j}{\sigma'(\rho)} + o(\rho^j), \tag{4}$$

which is sufficient for our purpose. Let $A_j = -1/\sigma'_j(\rho_j)$ be the residue of $1/(1 - \sigma_j(z))$ at $\rho_j.$ Then

$$\frac{1}{1 - \sigma_j(z)} - \frac{A_j}{z - \rho_j}$$

is analytic in the disc $\{z : |z| \leq \rho + \delta\}$ and by the Cauchy integral formula we get

$$a_{n,j} = -\frac{A_j}{\rho_j}(\rho_j)^{-n} + O\left(\frac{1}{(\rho + \delta)^n}\right).$$

Finally, since

$$\sigma_j(z) = \sigma(z) - \frac{z^j}{1 - z^j},$$

we get

$$\sigma'_j(\rho_j) = \sigma'(\rho_j) - \frac{j\rho_j^{j-1}}{(1 - \rho_j^j)^2}.$$

Hence, taking into account (4) we get

$$A_j = -\frac{1}{\sigma'(\rho)} + O(j\rho^j).$$

Consequently, for $j \geq 6$

$$a_{n,j} = \left(\frac{1}{\sigma'(\rho)} + O(j\rho^j)\right) \frac{1}{\left(\rho + \rho^j \frac{1+o(1)}{\sigma'(\rho)}\right)^{n+1}} + O((\rho + \delta)^{-n}), \tag{5}$$

for some $\delta > 0$ (universal for $j \geq 6$).

4. ASYMPTOTICS

The following claim will account for most of the asymptotic analysis of (1)

LEMMA 3. As $n \rightarrow \infty$,

$$\sum_{j=1}^n \left(1 - \frac{a_{n,j}}{a_n}\right) = \sum_{j=0}^{\infty} (1 - (1 - \rho^j \alpha)^n) + o(1),$$

where $\alpha = 1/\sigma'(\rho)$.

This statement immediately implies our theorem since the asymptotic behavior of the series on the right is known (see e.g., [2], [4], [7, Section 7.8 and references therein]).

PROOF OF LEMMA 3. The sequence (a_n) was studied in [5] and one has

$$a_n = \frac{1}{\sigma'(\rho)} \left(\frac{1}{\rho}\right)^{n+1} + O((\rho + \delta)^{-n}).$$

Combining this with (5), for $j \geq 6$

$$\frac{a_{n,j}}{a_n} = \frac{1 + O(j\rho^j)}{\left(1 + \rho^{j-1} \frac{1+o(1)}{\sigma'(\rho)}\right)^{n+1}} + O((1 + \Delta)^{-n}),$$

where $\Delta = \delta/\rho > 0$. Consequently,

$$\begin{aligned} \sum_{j=1}^n \left(1 - \frac{a_{n,j}}{a_n}\right) &= \sum_{j=1}^5 \left(1 - \frac{a_{n,j}}{a_n}\right) \\ &\quad + \sum_{j=6}^n \left(1 - \frac{1 + O(j\rho^j)}{\left(1 + \rho^{j-1} \frac{1+o(1)}{\sigma'(\rho)}\right)^{n+1}} + O((1 + \Delta)^{-n})\right). \end{aligned}$$

We will show that

$$\sum_{j=6}^n \frac{O(j\rho^j)}{(1 + \rho^{j-1} \alpha_j)^{n+1}} = o(1),$$

where we have set $\alpha_j = 1/\sigma'(\rho) + o(1)$. To this end, we will split this sum as

$$\left(\sum_{j=6}^{j_n} + \sum_{j=j_n+1}^n\right) \frac{O(j\rho^j)}{(1 + \rho^{j-1} \alpha_j)^{n+1}},$$

where j_n will be chosen momentarily. Since $\alpha_j \rightarrow 1/\sigma'(\rho) > 0$ we have

$$\sum_{j=j_n+1}^n \frac{O(j\rho^j)}{(1 + \rho^{j-1} \alpha_j)^{n+1}} \leq C \sum_{j>j_n} j\rho^j = O(j_n \rho^{j_n}) = o(1),$$

as long as $j_n \rightarrow \infty$ at any rate. The other term of the sum is bounded by

$$C \sum_{j=6}^{j_n} (1 + \rho^{j-1} \alpha_j)^{-n} \leq C j_n \exp(-n \ln(1 + \rho^{j_n-1} \alpha_{j_n})).$$

Since $\ln(1+x) \geq x - x^2/2$, for $x \geq 0$ we get the bound

$$Cj_n \exp\left(-n\alpha\rho^{j_n-1}\left(1 - \frac{\alpha\rho^{j_n-1}}{2}\right)\right) \leq Cj_n \exp(-cn\rho^{j_n-1}).$$

Choosing $j_n \sim \mu \log_{1/\rho} n$ we see that this expression is bounded by

$$O(\log n \cdot e^{-cn^{1-\mu}}) = o(1),$$

whenever $\mu < 1$.

It remains to consider the sum

$$\sum_{j=5}^{n-1} \left(1 - \left(1 - \frac{\rho^j \alpha_j}{1 + \rho^j \alpha_j}\right)^n\right).$$

We first replace this sum by a more convenient one

$$\sum_{j=5}^{n-1} (1 - (1 - \rho^j \alpha_j)^n).$$

The difference is at most

$$\sum_{j=5}^{\infty} \left(\left(1 - \frac{\rho^j \alpha_j}{1 + \rho^j \alpha_j}\right)^n - (1 - \rho^j \alpha_j)^n\right).$$

For a j_0 which will be chosen momentarily, we consider

$$\sum_{j=5}^{j_0} \left(\left(1 - \frac{\rho^j \alpha_j}{1 + \rho^j \alpha_j}\right)^n - (1 - \rho^j \alpha_j)^n\right).$$

Since each summand is nonnegative, this sum can be upperbounded by

$$\begin{aligned} \sum_{j=5}^{j_0} \left(1 - \frac{\rho^j \alpha_j}{1 + \rho^j \alpha_j}\right)^n &\leq \sum_{j=5}^{j_0} \exp\left(-\frac{\alpha_j \rho^j n}{1 + \rho^j \alpha_j}\right) \\ &\leq (j_0 + 1) \exp\left(-\frac{\alpha \rho^{j_0} n}{1 + \rho^{j_0} \alpha_j}\right) \leq (j_0 + 1) \exp(-c\rho^{j_0} n), \end{aligned}$$

for some absolute constant c . By choosing j_0 so that

$$c\rho^{j_0} n \geq \kappa \ln n,$$

i.e.,

$$j_0 \leq \frac{\ln((cn)/(\kappa \ln n))}{\ln(1/\rho)},$$

we see that the sum up to j_0 is bounded by $c \log n/n^\kappa$. For the remaining range of j 's we write

$$\begin{aligned} \left(1 - \frac{\rho^j \alpha_j}{1 + \rho^j \alpha_j}\right)^n - (1 - \rho^j \alpha_j)^n &= \left(1 - \rho^j \alpha_j + \frac{\rho^{2j} \alpha_j^2}{1 + \rho^j \alpha_j}\right)^n - (1 - \rho^j \alpha_j)^n \\ &\leq n \frac{\alpha^2 \rho^{2j}}{1 + \rho^j \alpha} \left(1 - \frac{\rho^j \alpha}{1 + \rho^j \alpha}\right)^{n-1} \leq n\alpha^2 \rho^{2j}, \end{aligned}$$

where we have used the inequality $(a + b)^n - a^n \leq nb(a + b)^{n-1}$ valid for nonnegative numbers a and b . Hence

$$\sum_{j \geq j_0} \left(\left(1 - \frac{\rho^j \alpha_j}{1 + \rho^j \alpha_j} \right)^n - (1 - \rho^j \alpha_j)^n \right) \leq n \alpha_j^2 \sum_{j \geq j_0} \rho^{2j} \leq cn \rho^{2j_0}.$$

The choice of j_0 so that $\rho^{j_0} = \Theta(\log n/n)$ is within the previous constraint, and for that choice we have

$$cn \rho^{2j_0} = \Theta\left(\frac{\log^2 n}{n}\right).$$

Using the same argument we can show that α_j can be replaced by its limit $\alpha = 1/\sigma'(\rho)$, i.e., that we have

$$\sum_{j=5}^{n-1} (1 - (1 - \rho^j \alpha_j)^n) = \sum_{j=5}^{n-1} (1 - (1 - \rho^j \alpha)^n) + o(1).$$

Finally, the sum on the right can be increased to

$$\sum_{j=0}^{\infty} (1 - (1 - \rho^j \alpha)^n),$$

since

$$\sum_{j \geq n} (1 - (1 - \rho^j \alpha)^n) \leq \alpha n \sum_{j \geq n} \rho^j = o(1)$$

and is thus negligible. Also, for each fixed j , with $1 \leq j \leq 5$, $a_{n,j}/a_n = o(1)$ as we will indicate in the appendix. This means that

$$\sum_{j=1}^5 \left(1 - \frac{a_{n,j}}{a_n} \right) = 5 + o(1),$$

and proves the Lemma. □

ACKNOWLEDGEMENTS

This paper owes its existence to the curiosity of Herbert Wilf, who brought the paper [5] to our attention by asking PH whether the methods used in [3] could be used to rederive some of the results of [5] (that question, by the way, remains unanswered).

REFERENCES

1. L. Carlitz, Restricted compositions, *The Fibonacci Quart.*, **14** (1976), 254–264.
2. P. Flajolet, X. Gourdon and P. Dumas, Mellin transform and asymptotics: harmonic sums. Special volume on mathematical analysis of algorithms, *Theor. Comput. Sci.*, **144** (1995), 3–58.
3. P. Hitczenko and C. D. Savage, On the multiplicity of parts in a random composition of a large integer, preprint (<http://www.mcs.drexel.edu/~phitczen>).
4. P. Hitczenko and G. Stengle, Expected number of distinct part sizes in a random integer composition, *Comb. Probab. Comput.*, **9** (2000), 519–527.
5. A. Knopfmacher and H. Prodinger, On Carlitz compositions, *Europ. J. Combinatorics*, **19** (1998), 579–589 (<http://www.idealibrary.com>).

6. G. Louchard and H. Prodinger, Probabilistic analysis of Carlitz compositions, preprint (<http://www.wits.ac.za/helmut/paper1st.htm>).
7. R. Sedgewick and P. Flajolet, *An Introduction to the Analysis of Algorithms*, Addison–Wesley, Reading, MA, 1996.
8. N. Sloane, *The On-line Encyclopedia of Integer Sequences*, (<http://www.research.att.com/~njas/sequences>).

APPENDIX

In order to show that $\sigma_j(z) = 1$ has a unique real root ρ_j and the existence of a $\delta > 0$ satisfying the asserted properties we rewrite $\sigma(z)$ in a more convenient form

$$\sigma(z) = \sum_{m=1}^{\infty} \frac{z^m}{1+z^m},$$

which can be done by expanding $z^j/(1-z^j)$ into geometric series and interchanging the order of summation. Now, $\sigma_j(z) = 1$ can be rewritten as

$$\sum_{m=1}^{\infty} \frac{z^m}{1+z^m} - \frac{z^j}{1-z^j} - 1 = 0.$$

We want to use Rouché's theorem. To this end split the left-hand side as $f(z) + g_j(z)$ where

$$f(z) = \sum_{m=1}^6 \frac{z^m}{1+z^m} - 1.$$

It can be verified that

$$\min_{|z|=0.663} |f(z)| \geq 0.28.$$

Actually Maple suggests that a stronger claim is true, namely that the minimum of $|f(z)|$ on that circle is attained at $z = 0.663$ and is 0.283467. We have not tried to prove it and thus we claim only the weaker statement, which can be verified by evaluating $|f|$ at sufficiently many points and using its Lipschitz property: for $|z_1| = |z_2| = r$,

$$\begin{aligned} ||f(z_1)| - |f(z_2)|| &\leq |f(z_1) - f(z_2)| \leq \sum_{m=1}^6 \frac{|z_1^m - z_2^m|}{|(1+z_1^m)(1+z_2^m)|} \\ &\leq |z_1 - z_2| \sum_{m=1}^6 \frac{mr^{m-1}}{(1-r^m)^2} \leq C|z_1 - z_2|, \end{aligned}$$

where $C \leq (1-r)^{-2} \sum_{m \geq 1} mr^{m-1} \leq (1-r)^{-4}$. Since for $|z| = 0.663$

$$\begin{aligned} |g_j(z)| &\leq \sum_{m=7}^{\infty} \frac{|z|^m}{1-|z|^m} + \frac{|z|^j}{1-|z|^j} \\ &\leq \frac{0.663^7}{(1-0.663^7)(1-0.663)} + \frac{0.663^6}{1-0.663^6} \leq 0.27, \end{aligned}$$

we see that

$$\min_{|z|=0.663} |f(z)| > \max_{|z|=0.663} |g_j(z)|,$$

uniformly for all $j \geq 6$. Thus $f + g_j$ and f have the same number of roots in the disc $|z| \leq 0.663$. But f has a unique (and thus necessarily real) root on that disc (this again may be verified by Maple after converting it to a polynomial equation). The existence of δ follows since the same argument can be repeated for a disc with a radius just a bit smaller. As for the monotonicity, it suffices to rewrite $\sigma_j(z) = 1$ as $\sigma(z) = 1/(1 - z^j)$ and notice that $\sigma(z)$ is increasing as a function of a real variable and that

$$\frac{1}{1 - z^{j_1}} < \frac{1}{1 - z^{j_2}},$$

whenever $j_1 > j_2$ and $0 < z < 1$ is real.

The same argument (with changed parameters) can be used to verify that for $j = 2, 3, 4$ and 5 , on the disc $|z| < r, 0 < r < 1, \sigma_j(z) = 1$ has a unique real root $\rho_j > \rho$. The hardest case is $j = 2$. We found that the splitting

$$f_2(z) = \sum_{m=1}^{25} \frac{z^m}{1 + z^m} - \frac{1}{1 - z^2}, \quad g(z) = \sum_{m=26}^{\infty} \frac{z^m}{1 + z^m},$$

will do the job. On the circle $|z| = 0.8$ one has $|f_2(z)| \geq 0.06, |g(z)| \leq 0.016$. The polynomial resulting from multiplying f_2 by the product of denominators is of degree 264 (after cancellations). Maple (somewhat reluctantly) shows that the root closest to zero is real and is about 0.78397 (there is another real root of about 0.927122) and the next closest to zero roots are complex conjugate and have absolute value around 0.81914. For $j = 3, 4, 5$ one can get away by letting $g(z) = \sum_{m=11}^{\infty} z^m / (1 + z^m)$, and choosing $r = 0.75$.

The same method could presumably be used to force the argument for $j = 1$. However, here matters would be computationally worse. Furthermore, the equation $\sigma_1(z) = 1$ does not have real solutions, and thus, its closest to zero root (if it exists) would have to come from a pair of complex conjugates. Luckily, for that case we can use a different argument based on the probabilistic approach used in [3] (see also [4]). Let us briefly sketch it. Consider the set C_n of all 2^{n-1} compositions of an integer n and let \mathbf{Q} be the uniform probability measure on C_n . Since the restriction of such a measure to any subset is again the uniform measure on that subset we can view the uniform measure \mathbf{P} on the set of all Carlitz compositions as a conditional measure obtained by restricting \mathbf{Q} to Ω_n , i.e.,

$$\mathbf{P}(\cdot) = \mathbf{Q}(\cdot | \Omega_n).$$

Let A_1^c be the set of all compositions of n that do not use part size 1 and recall that I_1^c is the set of all Carlitz compositions with this property. Then,

$$\frac{a_{n,1}}{a_n} = \mathbf{P}(I_1^c) = \mathbf{Q}(A_1^c | \Omega_n) = \frac{\mathbf{Q}(A_1^c \cap \Omega_n)}{\mathbf{Q}(\Omega_n)} \leq \frac{\mathbf{Q}(A_1^c)}{\mathbf{Q}(\Omega_n)}.$$

Now, by the result of Knopfmacher and Prodinger

$$\mathbf{Q}(\Omega_n) \geq c \frac{1.75^n}{2^n} = c(0.875)^n,$$

and we need to upper bound $\mathbf{Q}(A_1^c)$. To this end we will use the observation made in [3] or [4] that a random composition of n is distributed like

$$(\tilde{\Gamma}_1, \dots, \tilde{\Gamma}_\tau),$$

where, with (Γ_i) being a sequence of i.i.d. geometric random variable with parameter $1/2$, we let

$$\tau = \inf \left\{ k : \sum_{q=1}^k \Gamma_q \geq n \right\},$$

and

$$\tilde{\Gamma}_q = \begin{cases} \Gamma_q & \text{if } q < \tau, \\ n - \sum_{p=1}^{\tau-1} \Gamma_p & \text{if } q = \tau. \end{cases}$$

Furthermore, τ is distributed like 1 plus a binomial random variable with parameters $n - 1$ and $1/2$. Therefore,

$$\begin{aligned} \mathbf{Q}(A_1^c) &= \mathbf{Q} \left(\bigcap_{k=1}^{\tau} \{ \tilde{\Gamma}_k > 1 \} \right) \leq \mathbf{Q} \left(\bigcap_{k=1}^{\tau-1} \{ \Gamma_k > 1 \} \right) \\ &\leq \mathbf{Q}(\tau \leq k_0) + \mathbf{Q} \left(\bigcap_{k=1}^{\tau-1} \{ \Gamma_k > 1 \} \cap \{ \tau > k_0 \} \right) \\ &\leq \mathbf{Q}(\tau - \mathbf{E}\tau \leq k_0 - \mathbf{E}\tau) + \mathbf{Q} \left(\bigcap_{k=1}^{k_0} \{ \Gamma_k > 1 \} \right) \\ &\leq \mathbf{Q} \left(|\tau - \mathbf{E}\tau| > \frac{n+1}{2} - k_0 \right) + \left(1 - \frac{1}{2} \right)^{k_0} \\ &\leq 2 \exp \left(-\frac{((n+1)/2 - k_0)^2}{2(n-1)/4} \right) + \exp(-k_0 \ln 2) \\ &\leq 2 \exp \left(-2 \frac{\phi_n^2}{n-1} \right) + \exp(-k_0 \ln 2), \end{aligned}$$

where we have put $k_0 = (n + 1)/2 - \phi_n$ and $\phi_n = \alpha(n + 1)$, for some $0 < \alpha < 1/2$. Then we get

$$\mathbf{Q}(A_1^c) \leq C(e^{-2\alpha^2})^n + C(e^{-(1/2-\alpha)\ln 2})^n.$$

To be able to claim that $\mathbf{Q}(A_1^c)/\mathbf{Q}(\Omega_n)$ tends to zero, we need both of the inequalities

$$e^{-2\alpha^2} < 0.875 \quad \text{and} \quad e^{-(1/2-\alpha)\ln 2} < 0.875$$

to be satisfied simultaneously. But this can be achieved by choosing α to be any number subject to $0.26 < \alpha < 0.3$. This completes the argument.

Received 7 February 2000 and accepted 28 April 2000

WILLIAM M. Y. GOH AND PAWEŁ HITCZENKO
 Department of Mathematics and Computer Science,
 Drexel University,
 Philadelphia,
 PA 19104, U.S.A.
 E-mail: wgoh@mcs.drexel.edu; phitzen@mcs.drexel.edu